

# U.S. FISH AND WILDLIFE SERVICE SPECIES ASSESSMENT AND LISTING PRIORITY ASSIGNMENT FORM

**Scientific Name:**

Quadrula houstonensis

**Common Name:**

Smooth Pimpleback

**Lead region:**

Region 2 (Southwest Region)

**Information current as of:**

03/03/2015

**Status/Action**

☐ Funding provided for a proposed rule. Assessment not updated.

☐ Species Assessment - determined species did not meet the definition of the endangered or threatened under the Act and, therefore, was not elevated to the Candidate status.

☐ New Candidate

☒ Continuing Candidate

☐ Candidate Removal

☐ Taxon is more abundant or widespread than previously believed or not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status

☐ Taxon not subject to the degree of threats sufficient to warrant issuance of a proposed listing or continuance of candidate status due, in part or totally, to conservation efforts that remove or reduce the threats to the species

☐ Range is no longer a U.S. territory

☐ Insufficient information exists on biological vulnerability and threats to support listing

- ☐ Taxon mistakenly included in past notice of review
- ☐ Taxon does not meet the definition of "species"
- ☐ Taxon believed to be extinct
- ☐ Conservation efforts have removed or reduced threats
- ☐ More abundant than believed, diminished threats, or threats eliminated.

## Petition Information

☐ Non-Petitioned

☒ Petitioned - Date petition received: 10/15/2008

90-Day Positive: 12/15/2009

12 Month Positive: 10/06/2011

Did the Petition request a reclassification? **No**

### For Petitioned Candidate species:

Is the listing warranted (if yes, see summary threats below) **Yes**

To Date, has publication of the proposal to list been precluded by other higher priority listing? **Yes**

Explanation of why precluded:

We find that the immediate issuance of a proposed rule and timely promulgation of a final rule for this species has been precluded by higher priority listing actions (including candidate species with lower LPNs), for the preceding 12 months, and continues to be precluded. During the past 12 months, the majority of our entire national listing budget has been consumed by work on various listing actions to comply with court orders and court-approved settlement agreements; meeting statutory deadlines for petition findings or listing determinations; emergency listing evaluations and determinations; and essential litigation-related administrative and program management tasks. We will continue to monitor the status of this species as new information becomes available. This review will determine if a change in status is warranted, including the need to make prompt use of emergency listing procedures. For information on listing actions taken over the past 12 months, see the discussion of Progress on Revising the Lists, in the current CNOR which can be viewed on our Internet website (<http://endangered.fws.gov/>).

## Historical States/Territories/Countries of Occurrence:

- **States/US Territories:** Texas
- **US Counties:** County information not available
- **Countries:** Country information not available

## Current States/Counties/Territories/Countries of Occurrence:

- **States/US Territories:** Texas
- **US Counties:** Austin, TX, Brazoria, TX, Brazos, TX, Burleson, TX, Burnet, TX, Colorado, TX, Coryell, TX, Falls, TX, Fayette, TX, Fort Bend, TX, Grimes, TX, Hamilton, TX, Leon, TX, Limestone, TX, Madison, TX, Matagorda, TX, McLennan, TX, Milam, TX, Robertson, TX, San Saba, TX, Waller, TX, Washington, TX, Wharton, TX
- **Countries:** Country information not available

## Land Ownership:

All of the known Smooth pimpleback populations in the Colorado or Brazos River systems are adjacent to private land.

## Lead Region Contact:

DIV OF ENDNGRD SPECIES AND HAB CONSERV, Nathan Allan, 512-490-0057,  
nathan\_allan@fws.gov

## Lead Field Office Contact:

## Biological Information

### Species Description:

The smooth pimpleback is a nearly round, thick-shelled freshwater mussel that generally reaches at least 60 mm (2.6 in) in length (Howells 2010b, p. 4). It is moderately thick, solid, and inflated. Externally, the smooth pimpleback, like its name suggests, is relatively smooth with minute sculpturing; it may or may not have a few small pustules (raised bumps) (Howells 2010b, p. 2). The external coloration of the shell ranges from tan to light brown, dark brown, and black with no rays (Howells 2010b, p. 4).

### Taxonomy:

The smooth pimpleback was originally described by Lea in 1859 as *Unio houstonensis*. It was later placed in the genus *Margarona* and ultimately moved to *Quadrula* by Simpson (1900, p. 782). Graf and Cummings (2007, p. 18) have proposed moving it to the genus *Amphinaia*, but other freshwater mussel taxonomists recommend waiting for additional work to be completed on

members of *Quadrula* before splitting the genus (Bogan 2011, pers. comm.). The smooth pimpleback is recognized by the Committee on Scientific and Vernacular Names of Mollusks of the Council of Systematic Malacologists, American Malacological Union (Turgeon et al. 1998, p 37), and we recognize it as a valid species.

## **Habitat/Life History:**

The smooth pimpleback has been found in mixed mud, sand, and fine gravel substrate in medium-to-large rivers and some reservoirs (Howells 2010b, p. 3) throughout its range. Currently, there is no specific information on age, or size of maturity. However, new research conducted by USGS and the Service on host fish use for central Texas Candidate mussels revealed that smooth pimpleback utilizes channel catfish as a host fish (Johnson et al. 2014, p. 43). Numerous individuals were examined for gravidity between June and November, with no evidence of eggs or glochidia (Howells 2000b, p. 38). Additionally, mussels in the genus *Quadrula* are typically short-term brooders (Gorden and Layzer 1989, p. 6; Garner et al. 1999, p. 277), and we expect the same of the smooth pimpleback. This species is a host fish specialist that freely broadcasts its glochidia by reflexive release in the form of mature conglomerates which are usually fragile and tend to disintegrate shortly after release (Johnson et al. 2014, p. 43).

Adult freshwater mussels are filter-feeders, siphoning algae, bacteria, detritus, microscopic animals, and dissolved organic matter (Fuller 1974, pp. 221-222, Silverman et al. 1997, p. 1862; Nichols and Garling 2000, p. 874-876; Christian et al. 2004, p. 109). For their first several months, juvenile mussels feed using cilia (fine hairs) on the foot to capture suspended as well as depositional material, such as algae and detritus (Yeager et al. 1994, pp. 253-259). Mussels tend to grow relatively rapidly for the first few years, and then slow appreciably at sexual maturity, when energy presumably is being diverted from growth to reproductive activities (Baird 2000, pp. 66-67). Mussels are extremely long lived, living from two to several decades (Rogers et al. 2001, p. 592), and possibly up to 200 years in extreme instances (Bauer 1992, p. 427).

Most mussel species, including smooth pimpleback, have distinct forms of male and female. During reproduction, males release sperm into the water column, which females draw in through their siphons. Fertilization takes place internally, and the resulting eggs develop into specialized larvae (called glochidia) within the female's modified gill pouch (called marsupia) for four to six weeks. The females will then release matured glochidia individually, in small groups, or embedded in larger mucus structures called conglomerates. Glochidia are obligate parasites (cannot live independently of their hosts) on fish and attach to the gills or fins of appropriate host species where they encyst (enclose in a cyst-like structure) and feed off of the host's body fluids (Vaughn and Taylor 1999, p. 913) and develop into juvenile mussels weeks or months after attachment (Arey 1932, pp. 214-215). The glochidia will die if they fail to find the appropriate host fish, attach to a fish that has developed immunity from prior infestations, or attach to the wrong location on a host fish (Neves 1991, p. 254; Bogan 1993, p. 299). Mussels experience their primary opportunity for dispersal and movement within the stream as glochidia attached to a host fish (Smith 1985, p. 105). Upon release from the host, newly transformed juveniles drop to the substrate on the bottom of the stream. Those juveniles that drop in unsuitable substrates die because their immobility prevents them from

relocating to more favorable habitat. Juvenile freshwater mussels burrow into interstitial substrates and grow to a larger size that is less susceptible to predation and displacement from high flow events (Yeager et al. 1994, p. 220). Throughout the rest of their life cycle, mussels generally remain within the same small area where they released from the host fish.

## **Historical Range/Distribution:**

The smooth pimpleback is native to the Brazos and Colorado River basins of central Texas (Karatayev and Burlakova 2008, p. 39). In the Colorado River basin, the species historically occurred throughout the length of the mainstem Colorado River from Coleman County downstream to Wharton County, as well as in the Llano River, and Onion and Skull Creeks. (Howells 1996, p. 21; 1997a, pp. 34–35; Randklev et al. 2010c, p. 4; OSUM 2011c, p. 1). Within the Brazos River basin, the species historically occurred throughout the length of the mainstem of the Brazos River (Howells 2009, p. 12), as well as in the Clear Fork Brazos, Leon, Navasota, Little Brazos, San Gabriel, Lampasas, and Little Rivers and Yegua Creek (Howells 2010b, pp. 4– 6; Randklev et al. 2010b, p. 20). The smooth pimpleback was historically uncommon where it occurred; from the 1960s through the 1990s, experts failed to find large populations persisting throughout its range (Howells 2009, p. 12).

The smooth pimpleback has also been reported from the Trinity River and other drainages in Texas, as well as from areas outside of Texas, including southern Arkansas and the Verdigris River in Kansas. These reports are likely misidentifications of other pimpleback species that can sometimes closely resemble smooth pimpleback (Howells 2010b, pp. 4-5).

## **Current Range Distribution:**

The smooth pimpleback has been nearly extirpated from the Colorado River basin. A few small populations persist in the Brazos River basin. Recent surveys suggest a greater abundance and distribution of the smooth pimpleback in the central and lower Brazos River drainage than was indicated by collections from the past 40 years, with five populations represented by more than a few individuals. Currently the smooth pimpleback is restricted to one mainstem reservoir, 11 sites in eight counties on the mainstem Colorado River, and throughout the lower San Saba River. Populations in all of the other historically occupied tributaries and two reservoirs appear to have been extirpated.

### **Colorado River System**

Between 1993 and 1999 numerous surveys in many locations on the Colorado River occurred with no evidence of smooth pimpleback, except for in Colorado County (Howells 1995, p. 29; 1996, p. 23; 1997a, pp. 27, 31; 2000a, p. 27; 2002a, p. 6; 2004, p. 7, 11; 2005, p. 6). It was not until after 1999 that the smooth pimpleback was found in the Colorado River in areas other than Colorado County. Historically, smooth pimpleback was found live within the mainstem of Colorado River in Coleman County; however, no surveys have been conducted in that area since 1989 to determine

whether or not it persists in that section of the river. In 2011, several live individuals were located in the mainstem Colorado River downstream from its confluence with the San Saba River in San Saba County (Randklev 2012, pers. comm.). Several very recently dead shells were located further downstream in San Saba and Lampasas Counties during presence/absence surveys in 2012 and 2013 (Service Files, 2012; Service Files, 2013).

Several surveys have taken place over the years within a series of three in-channel reservoirs on the Colorado River that fall within Burnet and Llano Counties. In 2001, one live smooth pimpleback was found in Lake Lyndon B. Johnson, a large mainstem reservoir in Burnet/Llano County, but no live individuals have been found since (Howells 2002a, pp. 6–7; 2006, pp. 68–69). Farther downstream in Inks Lake, a small mainstem reservoir in the same counties, several live smooth pimpleback were found in 1992 (Howells 1994, p. 4); however, since that time only shell material has been found during four separate surveys between 1996 and 2005 (Howells 1997a, pp. 32–33; 1999, p. 16; 2005, p. 8; 2006, p. 67). Frequent drawdowns in this lake appear to have affected all species of freshwater mussels, as there has been a sharp decline in the overall mussel community (Howells 1999, p. 16). Farther downstream in Lake Marble Falls, another small mainstem reservoir in Burnet County, 13 live and several recently dead smooth pimpleback were found in 1995 during a drawdown of lake levels (Howells 1996, p. 22), but subsequent surveys in 1996 failed to find any additional living animals (Howells 1997a, p. 33). The small recent survey effort is not sufficient to conclude that the smooth pimpleback no longer occur in these lakes, and small populations may still persist there.

In 1999, three live individuals were located within the mainstem of the Colorado River in Colorado County (Howells 2000a, p. 27). Live smooth pimpleback were found in the same general area during two surveys in 2009 (Burlakova and Karatayev 2010a, p. 16; Johnson 2009, p. 1). Presence/absence surveys in 2012 at this site and farther downstream found smooth pimpleback to be a dominate species in these areas. In addition, a large population was discovered partially exposed during low water conditions and relocated to deeper water (Service Files, 2012; Service Files, 2013). The Service also conducted surveys at another site in Colorado County and another site in Wharton County in 2013 and only found very recently dead to long dead individuals (Service Files, 2013). Farther downstream, at a different site in Wharton County, live smooth pimpleback were found at two sites in 2009 (Burlakova and Karatayev 2010a, p. 16), despite having been surveyed in 1995 where none were found (Howells 1996, p. 23).

Smooth pimpleback were found at two locations in the San Saba River in San Saba County, when 29 individuals were found at two locations in 2011 (Burlakova and Karatayev 2011, p. 5; Randklev 2011b, p. 1). Various size and age classes were represented, indicating a reproducing, recruiting population in the riffle and pool habitats (Burlakova and Karatayev 2011, p. 5). Even more recently, 206 smooth pimpleback, including adults and juveniles, were recorded in this same area in riffle and pool habitat (Randklev 2011b, p. 1). The Service conducted presence/absence surveys during 2012 at several sites throughout San Saba County and found several live smooth pimpleback of various age classes (Service Files, 2012). The population size of smooth pimpleback in the San Saba River may be approximately 232,920 mussels (Burlakova and Karatayev 2012b, p. 13).

No known extant populations of smooth pimpleback have been identified in any of the other Colorado River tributaries in which the species was historically known to occur, including the full length of the Llano River (Howells 1996, pp. 21–22; 1998, p. 17; 2000a, p. 25; 2005, p. 8; Randklev et al. 2010c, p. 4; OSUM 2011c, p. 1). A single subfossil shell, likely a smooth pimpleback, was found in the Llano River in Kimble County in 1995 (Howells 1996, pp. 21–22), but no other evidence of the species has been found throughout the Llano and Pedernales Rivers in recent years (Service Files, 2012). Additionally, although Onion and Skull Creeks were historically occupied by smooth pimpleback (Randklev et al. 2010c, p. 4), the species has not been found recently in either stream (Howells 1995, pp. 28–29; Service Files, 2012).

## Brazos River System

The smooth pimpleback historically occurred in the Brazos River system from Haskell and Throckmorton Counties downstream to Austin and Waller Counties, as well as in numerous tributaries. The species has been extirpated from the upstream half of the mainstem Brazos River and from at least three tributaries. Substantial populations persist in the Leon River, Navasota River, and Yegua Creek, and small populations remain in the lower Brazos and Little Brazos Rivers.

In the upper mainstem Brazos River, surveys in Palo Pinto, Somervell, and Bosque Counties between 1996 and 2000 indicate that the smooth pimpleback has been extirpated from the upstream portion of the river (Howells 1997a, pp. 16, 18–19; 1999, pp. 11–12; 2001, p. 19).

Although not extirpated from the middle Brazos River, the smooth pimpleback occurs only in low numbers. Surveys in 1996 and 1998 found no individuals (Howells 1997a, p. 21; 1999, p. 12). A single live smooth pimpleback was found in McLennan County in the middle Brazos River in 2005 (Howells 2010b, p. 5). Two live and several recently dead individuals were recorded in Falls County in 2006 (Karatayev and Burlakova 2008, pp. 6–10). In Milam and Robertson Counties, no smooth pimpleback were found in 1998 (Howells 1999, p. 13), but eight live individuals were found in 2006 (Burlakova and Karatayev 2010b, p. 1). More recently, in 2008, 13 live smooth pimpleback were found at the same site (Randklev et al. 2009, p. 18). Additionally, downstream in Burleson and Brazos Counties, which were historically occupied by the smooth pimpleback (OSUM 2011c, p. 1), a small population persists. In 1995, one live and one recently dead individual were collected within Brazos County (Howells 1996, pp. 17–18). Although none were found here in 1999 (Howells 2000a, pp. 21–22), in 2006 a single live smooth pimpleback was collected at this site (Karatayev and Burlakova 2008, pp. 6–10). Surveys conducted at three different locations in Brazos and Washington Counties in 2012 found a total of 22 individuals (Randklev 2012, pers. comm.). In 2014, the area was revisited by the Service and smooth pimpleback was confirmed to continue to persist and is successfully recruiting (Service Files, 2014). Additionally, further downstream in Grimes and Waller Counties, a single live individual was found in 2006 (Burlakova and Karatayev 2010b, p. 1) and again in 2008 (Randklev et al. 2009, p. 18).

Smooth pimpleback are more numerous in the lower mainstem Brazos River. Thirty-eight live were found in Austin and Waller Counties in 2006 (Karatayev and Burlakova 2008, pp. 6–10), and 59 live

individuals were found there in 2012 (Randklev 2012, pers. comm.). Farther downstream in Austin and Fort Bend Counties only two live smooth pimpleback were found in 2012 (Randklev 2012, pers. comm.). Surveys in Fort Bend County in 2012, found 25 live individuals (Randklev 2012, pers. comm.). There are no historical records of this species occurring as far downstream as Fort Bend County.

Historically, the smooth pimpleback occurred in numerous tributaries to the Brazos River but now can only be found in sections of Leon, Little, Navasota, and Little Brazos Rivers and Yegua Creek. Only subfossil shells of smooth pimpleback have been found in the Clear Fork Brazos River (Howells 1999, p. 19) since it was originally documented in Shackelford County in 1893 (Randklev et al. 2010c, p. 4). Surveys conducted during 2009 to 2011 in Haskell, Throckmorton, Jones, and Shackelford Counties found subfossil shells (Cedar Ridge Reservoir Project-Draft 2011) indicating that the smooth pimpleback once occurred throughout the Clear Fork Brazos River system. This species has most likely been extirpated from this system considering no live or recently dead smooth pimpleback have been documented.

The Leon River, in the Little River drainage of the Brazos, historically contained smooth pimpleback throughout its length in Comanche, Hamilton, Coryell, and Bell Counties (Howells 1994, p. 19, 1997a, p. 20; Randklev et al. 2010c, p. 4; OSUM 2011c, p. 1). Currently, a smooth pimpleback population persists in Hamilton County, where numerous live individuals were found in 2006 and 2011 (Howells 2006, pp. 82–83; Randklev 2011a, p. 1), as well as several locations in Coryell County, where numerous individuals were found in 2011 (Randklev 2011a, p. 1). A few live individuals were located in Comanche and Bell Counties as well in 2011 (Randklev 2011a, p. 4).

Only subfossil smooth pimpleback shells have been found in the Lampasas River, also in the Little River drainage, in Bell County in 1996 (Howells 1997a, pp. 20, 23). Subsequent surveys of the river in both Bell and Lampasas Counties yielded no evidence of smooth pimpleback (Howells 1999, p. 14; 2001, p. 20), and the species has likely been extirpated from the Lampasas River.

The Little River in Milam County is also a historical location for the smooth pimpleback (Randklev et al. 2010c, p. 4). Old shells were found at this site in 1996 (Howells 1997a, p. 22), and a single live individual was found here in 2006 (Karatayev and Burlakova 2008, p. 6). Because smooth pimpleback populations have been located above and below the confluence of Little River in the mainstem Brazos River, there is a possibility that these populations could influence recruitment within the Little River. Based on this information, the Service believes that a small population may continue to persist in the mainstem Little River in Milam County but in very low numbers.

A single old smooth pimpleback shell has been found in the San Gabriel River in Milam County (Howells 1997a, p. 23). However, no recent surveys have been conducted to determine the current status of the smooth pimpleback in this river.

In the Little Brazos River, the smooth pimpleback appears to persist in low numbers. Although none were found in Robertson County in 1993 and there had appeared to be a die off of numerous freshwater mussel species (Howells 1995, p. 18), one live smooth pimpleback was found during a 2006 survey (Karatayev and Burlakova 2008, p. 6). Farther downstream in Brazos County, recently



dead individuals were discovered in 2001 (Howells 2002a, pp. 4–5). The species occurred in this area historically (Randklev et al. 2010c, p. 4), and reports of mussels in the Little Brazos River from the 1950s described the freshwater mussel community as numerous, including smooth pimpleback (Gentner and Hopkins 1966, pp. 458–459). If smooth pimpleback persists in this river, it is most likely in low numbers.

In the Navasota River, smooth pimpleback historically occurred in Leon, Brazos, Grimes, and Washington Counties (Randklev et al. 2010c, p. 4; OSUM 2011c, p. 1). Currently, the species persists in each of those counties, with a large population occurring in the lower river. In Leon County three recently dead smooth pimpleback shells were found in 2000 (Howells 2001, p. 23), indicating that a few individuals may persist in the area. However, one of the largest known populations occurs farther downstream near the confluence of the Navasota and Brazos Rivers. Nine live individuals were found in this area in 2006 (Karatayev and Burlakova 2008, pp. 6–10), and in 2008 a total of 117 live smooth pimpleback were recorded at 3 different locations within Washington and Grimes Counties (Randklev et al. 2009, pp. 6, 18). A large population continues to persist in the Navasota River, with a total of 314 smooth pimpleback recorded at two sites in 2011 (Randklev 2011a, p. 1).

In Yegua Creek, subfossil smooth pimpleback shells were found in Washington County in 1996. No smooth pimpleback were found during several surveys between 1996 and 2003 (Howells 1997a, pp. 24–26; 2001, p. 22; 2004, p. 6). However, in 2006, a live individual was discovered (Karatayev and Burlakova 2008, pp. 6–10), which prompted further surveys in 2008. Numerous smooth pimpleback were found during subsequent surveys at four different locations within Washington and Burleson Counties (Randklev et al. 2009, pp. 16–18; Randklev 2011a, p. 1), indicating the presence of a potentially large population in this stream.

## **Population Estimates/Status:**

Based on historical and current data, the smooth pimpleback has declined rangewide and is now known from only nine counties throughout the Colorado River basin and it occurs in 14 counties throughout the Brazos River basin. The species has been eliminated from nearly the entire Colorado River and all but one of its tributaries, and has been eliminated from the upper Brazos River and several tributaries as well. The lower Colorado River, San Saba River, lower Brazos River, Navasota River, Leon River, and Yegua Creek populations appear to be stable and reproducing, but the remaining populations are small, isolated, and represented by only a few individuals.

## **Threats**

### **A. The present or threatened destruction, modification, or curtailment of its habitat or range:**

The decline of mussels in Texas and across the United States is primarily the result of habitat loss and degradation (Neves 1991, pp. 252, 265; Howells et al. 1996, pp. 21–22). Chief among the causes of mussel decline in Texas are the effects of impoundments, sedimentation, dewatering, sand and gravel mining, and chemical contaminants (Neck 1982a, pp. 33–35; Howells et al. 1996, pp. 21–22; Winemiller et al. pp. 17–18). These threats are discussed below.

## Impoundments

A major factor in the decline of freshwater mussels across the United States has been the large-scale impoundment of rivers (Vaughn and Taylor 1999, p. 913). Dams are the source of numerous threats to freshwater mussels: They block upstream and downstream movement of species by blocking host fish movement; they eliminate or reduce river flow within impounded areas, thereby trapping silts and causing sediment deposition; and dams change downstream water flow timing and temperature, decrease habitat heterogeneity, and affect normal flood patterns (Layzer et al. 1993, pp. 68–69; Neves et al. 1997, pp. 63–64; Watters 2000, pp. 261–264; Watters 1996, p. 80). Within reservoirs (the impounded waters behind dams), the decline of freshwater mussels has been attributed to sedimentation, decreased dissolved oxygen, and alteration of resident fish populations (Neves et al. 1997, pp. 63–64; Pringle et al. 2000, pp. 810–815; Watters 2000, pp. 261–264). Dams significantly alter downstream water quality and stream habitats (Allan and Flecker 1993, p. 36; Collier et al. 1996, pp. 1, 7) resulting in negative effects to tailwater (the area downstream of a dam) mussel populations (Layzer et al. 1993, p. 69; Neves et al. 1997, p. 63; Watters 2000, pp. 265–266). Below dams, mussel declines are associated with changes and fluctuation in flow regime, scouring and erosion of stream channels, reduced dissolved oxygen levels and water temperatures, and changes in resident fish assemblages (Williams et al. 1992, p. 7; Layzer et al. 1993, p. 69; Neves et al. 1997, pp. 63–64; Pringle et al. 2000, pp. 810–815; Watters 2000, pp. 265–266). Numerous dams have been constructed throughout the Colorado and Brazos River systems within the range of smooth pimpleback (Stanley et al. 1990, p. 61).

Population losses due to the effects of dams and impoundments have likely contributed more to the loss of diversity and abundance of freshwater mussels across Texas, including the smooth pimpleback, than any other factor. Stream habitat throughout nearly all of the range of smooth pimpleback has been affected by numerous impoundments, leaving generally short, isolated patches of remnant habitat between dams. Impoundments have resulted in profound changes to the nature of the rivers, primarily replacing free-flowing river systems with a series of large reservoirs.

There are no natural lakes within the range of the smooth pimpleback, however, the species is able to tolerate some impoundment conditions and have been known to occur in three mainstem reservoirs on the Colorado River, although all but one population is likely extirpated (Howells 1997a, pp. 32–33; 1999, p. 16; 2005, p. 8; 2006, p. 67). Dams continue to fragment smooth pimpleback populations, and the downstream effects of dams are detrimental to their habitat.

Impoundments occur throughout the range of the smooth pimpleback. The majority of Nueces-Frio, Guadalupe-San Antonio, Colorado, and Brazos Rivers, as well as many tributaries, are now

impounded. There are 74 major reservoirs and numerous smaller impoundments within the historical and current range of the smooth pimpleback. Thirty-one of the 74 major reservoirs are located within the Colorado River basin, and the remaining 43 reservoirs are located within the Brazos River basin. There are also eleven new reservoirs that have been recommended for development as feasible alternatives to meet future water needs within the Brazos River basin (Brazos G Regional Water Planning Group 2010, p. 4B.12–1). In addition, six new off-channel reservoirs are also being considered for future development (Brazos G Regional Water Planning Group 2010, p. 4B.13–2). At least one of the proposed reservoir sites on the Little River in Milam County is in the vicinity of where a single live smooth pimpleback was found in 2006 (Karatayev and Burlakova 2008, p. 6). These, and numerous smaller dams, occur throughout the Colorado and Brazos River basins and have resulted in ongoing destruction and modification of smooth pimpleback habitat and the curtailment of its range.

Dams threaten freshwater mussels in several ways. First, they can prevent the movement of freshwater mussel host fish. The overall distribution of mussels is a function of the dispersal of their hosts (Watters 1996, p. 83). For example, Watters (1996, p. 80) found that the distributions of the fragile papershell (*Leptodea fragilis*) and pink heelsplitter (*Potamilus alatus*) in five midwestern rivers were determined by the presence of low-head dams. These dams were non-navigable (without locks), lacked fish ladders, and varied in height from 1 to 17.7 m (3 ft to 58 ft), and the host fish could not disperse through them. Although the distribution of mussels may depend on many ecological factors, the evidence presented in Watters (1996, pp. 79–85) illustrates that dams as small as 1 m (3 ft) high can limit the distribution of mussels. There are many dams that occur throughout the range of the smooth pimpleback that lack fish ladders and may be a barrier to the movement of fish hosts and, therefore, the distribution of mussels. Because the smooth pimpleback populations are all separated by dams of various sizes that are not passable by fish, the mussel is unable to disperse from its current occupied range through host fish migration.

Dams also alter aquatic habitat within the resulting impoundments. It is well documented that many mussel species that are adapted to flowing water stream environments do poorly in the altered aquatic conditions found within impoundments (Williams et al. 1992, p. 7; Vaughn and Taylor 1999, p. 913). Once a dam is constructed, the original river channel upstream remains intact but under much deeper water with much lower velocities. As water velocity decreases, water loses its ability to carry sediment; sediment falls to the substrate, eventually smothering mussels that cannot adapt to soft substrates (Watters 2000, p. 263). Over time, the original mussel species composition of the stream channel may be eliminated or changed in favor of silt tolerant species (Watters 2000, p. 264). The mussel community may be altered from one with many different species to a community dominated by one to several very common species (Neck 1982b, p. 174). For species such as smooth pimpleback that may be able to survive the initial inundation of reservoirs, conditions within the reservoir are likely to become uninhabitable. The deep water in reservoirs is very cold and often devoid of oxygen and necessary nutrients (Watters 2000, p. 264). Cold water (less than 11 ° C (52 °F)) has been shown to stunt mussel growth (Hanson et al. 1988, p. 352). Because mussel reproduction is temperature dependent (Watters and O'Dee 1999, p. 455), it is likely that individuals living in the constantly cold hypolimnion in these channels may never reproduce, or reproduce less frequently (Watters 2000, p. 264). Any smooth pimpleback that survived the initial inundation may

have been unable to reproduce, eventually eliminating the species from large areas of the reservoir. The same would be true for mussels living in coldwater discharges downstream of large impoundments (Watters 2000, p. 264). The inundation of stream habitat by impoundments is a likely cause of the reduction in the distribution of the smooth pimpleback. The presence of the impoundments has caused the permanent loss of smooth pimpleback habitat throughout its range.

Mussels downstream of impoundments are often affected through changes in fish host availability, water quality (particularly lower water temperatures), habitat structure, and stream channel scouring (Vaughn and Taylor 1999, p. 916). The release of cold water from the hypolimnion (deeper and colder layer of water in reservoirs) can decrease the occurrence of fish species adapted to warm water and increase the occurrence of fish species adapted to colder water (Edwards 1978, pp. 73–75). This changes the species composition of suitable host fish and may prevent mussels from completing an essential part of their reproductive cycle. This has been demonstrated by the extirpation of mussel species from several rivers on the eastern seaboard of the United States, which has been linked to the disappearance of appropriate host fish; the reintroduction of the host fish to rivers has enabled mussel species to recolonize areas (Kat and Davis 1984, p. 174). In addition, because mussel reproduction is temperature dependent (Watters and O'Dee 1999, pp. 455–456), it is likely that individual mussels living in cold waters downstream of dam releases may reproduce less frequently, if at all (Layzer et al. 1993, p. 69). Low water temperatures can also significantly delay or prevent metamorphosis and glochidial release, which is often triggered by water temperature (Watters and O'Dee 1999, pp. 454–455; Watters and O'Dee 2000, p. 136).

In addition to the temperature of water released from dams, highly fluctuating, turbulent tailwaters devoid of sediment will scour the riverbed downstream of dams, rendering the area without mussel habitat (Layzer et al. 1993, p. 69). Depending on the use of the dam, water levels may fluctuate on a regular interval (for hydroelectric purposes) or at random (for flood control) (Watters 2000, p. 265). On the Colorado River, Inks Lake, Lake Marble Falls, Lake Buchanan, Lake Austin, Lake Travis, and Lady Bird Lake are each used for one or both of these purposes. Mortality of another rare mussel species in Texas, the Texas heelsplitter (*Potamilus amphichaenus*) was attributed to scheduled dewatering of the Neches River below B.A. Steinhagen Reservoir in east Texas (Neck and Howells 1994, p. 15).

In one study of the downstream effects of dams, Vaughn and Taylor (1999, p. 915) found a strong, gradual, linear increase in mussel species richness and abundance at sites on the Little River in Oklahoma downstream from Pine Creek Reservoir. Their research revealed that mussel species richness and total abundance did not begin to rebound until 20 km (12 mi) downstream of the impoundment and did not peak until 53 km (33 mi) downstream. They noted the most obvious difference since reservoir construction has been the alteration of the flow and temperature regimes, which gradually return to preimpoundment levels with downstream distance from the dam. These alterations appear to have produced an extinction gradient of mussels that is most severe near the dam (Vaughn and Taylor 1999, p. 915). We expect similar effects of the smooth pimpleback and

other Texas mussels downstream of dams. For example, mussel habitat below the Possum Kingdom Reservoir in the mainstem of the Brazos River did not stabilize until 150 km (240 mi) below the dam (Yeager 1993, p. 68).

Dam construction also fragments the range of smooth pimpleback, leaving remaining habitats and populations isolated by the structures as well as by extensive areas of deep, uninhabitable, impounded waters. These isolated populations are unable to naturally recolonize suitable habitat that may be impacted by temporary but devastating events, such as severe drought, chemical spills, or unauthorized discharges. Dams impound river habitats throughout almost the entire range of the species. These impoundments have left short and isolated patches of remnant habitat, typically in between impounded reaches.

The widespread construction of dams throughout the range of smooth pimpleback has significantly altered stream habitat both upstream and downstream of the dams by changing fish assemblages, temperature, dissolved oxygen, and substrate. The effects of dams are ongoing, decades after construction. In addition, the construction of new reservoirs is also being considered within the species' range that could result in additional habitat loss. Because of this loss of habitat and its effects on the populations, we conclude that the effects of impoundments are a threat to the smooth pimpleback.

## Sedimentation

Siltation and general sediment runoff is a pervasive problem in streams and has been implicated in the decline of stream mussel populations (Ellis 1936, pp. 39–40; Vannote and Minshall 1982, p. 4105; Dennis 1984, p. ii; Brim Box and Mossa 1999, p. 99; Fraley and Ahlstedt 2000, pp. 193–194). Specific biological effects on mussels from excessive sediment include reduced feeding and respiratory efficiency from clogged gills (Ellis 1936, p. 40), disrupted metabolic processes, reduced growth rates, increased substrate instability, limited burrowing activity (Marking and Bills 1979, pp. 208–209; Vannote and Minshall 1982, p. 4106), physical smothering, and disrupted host fish attractant mechanisms (Hartfield and Hartfield 1996, p. 373). The primary effects of excess sediment on mussels are sublethal, with detrimental effects not immediately apparent (Brim Box and Mossa 1999, p. 101).

The physical effects of sediment on mussel habitats are multifold and include changes in suspended material load; changes in streambed sediment composition from increased sediment production and runoff in the watershed; changes in the form, position, and stability of stream channels; changes in water depth or the width-to-depth ratio, which affects light penetration and flow regime; actively aggrading (filling) or degrading (scouring) channels; and changes in channel position that may leave mussels stranded (Brim Box and Mossa 1999, pp. 109–112). Increased sedimentation and siltation may explain, in part, why smooth pimpleback appear to be experiencing recruitment failure in some streams. Interstitial spaces (small openings between rocks and gravels) in the substrate provide essential habitat for juvenile mussels. When clogged with sand or silt, interstitial flow rates and spaces may become reduced (Brim Box and Mossa 1999, p. 100), thus reducing juvenile habitat availability. Juvenile freshwater mussels, including smooth pimpleback

juveniles, burrow into interstitial substrates, making it particularly susceptible to degradation of this habitat.

Even in 1959, Colorado River was noted as having high sedimentation rates from agricultural activities (Soil Conservation Service 1959, pp. 56, 59). Approximately 40 percent of U.S. river miles do not meet Clean Water Act standards due to excessive sediment loads (Environmental Protection Agency (EPA) 2000, p. 1), with agricultural activities being the primary source of sediment in streams (Waters 1995, p. 170). In general, sedimentation, resulting from unrestricted access by livestock, has been shown to be a significant threat to many streams and their mussel populations (Fraley and Ahlstedt 2000, p. 193).

The dominant land use in the Colorado River basin is grazing (Hersh 2007, p. 11). Soil compaction from intensive grazing may reduce infiltration rates and increase runoff, and trampling of riparian vegetation increases the probability of erosion (Armour et al. 1994, p. 10; Brim Box and Mossa 1999, p. 103). Additionally, much of the Brazos River basin is grazed or farmed for row crops, which often contributes large amounts of sediment to the basin (Brazos River Authority 2007, p. 4). Reservoir construction in the upper portion of the basin has been attributed with the erosion and subsequent sedimentation of the lower river (USGS 2001, p. 30), as sediment-poor tailwaters scour the riverbanks below the dam and deposit sediment farther downstream. In 2004, sedimentation was high enough in the Brazos River below Possum Kingdom Reservoir to cause residents to raise concerns to the Brazos River Authority (Brazos River Authority 2006, p. 2), and elevated suspended sediment levels have been reported throughout the basin (Brazos River Authority 2006, p. 8).

Sedimentation may become an increasing threat to the smooth pimpleback in the Colorado and Brazos River basins as the Austin metropolitan area continues to expand. Activities associated with urbanization, such as road construction, increased impervious surfaces, and road construction can be detrimental to stream habitats (Couch and Hamilton 2002, p. 1). The City of Austin, population approximately 800,000 people (Austin City Connection 2011, p. 1) lies within the Colorado River basin, and 3.9 million people live within the Brazos River basin (Brazos River Authority 2007, p. 1). Both of these basins have undergone substantial urbanization providing sources of increased sediment runoff into habitats of the smooth pimpleback. Runoff from increased impervious surfaces increases sediment loads in streams and destabilizes stream channels (Pappas et al. 2008, p. 151). Impervious surfaces also result in channel instability by accelerating stormwater runoff, which increases bank erosion and bed scouring, thereby further increasing downstream sedimentation (Brim Box and Mossa 1999, p. 103). While erosion and sedimentation associated with road construction may be temporary, the existence of road crossings is shown to have ongoing impacts to mussel habitat. For example, in the Guadalupe River, road crossings were found to cause a long-term increase in sedimentation both upstream and downstream, as channel constriction reduced flow upstream, causing sediment deposition, and runoff from the road increased sedimentation downstream (Keen- Zebert and Curran 2009, p. 301). Urban development activities may also affect streams and their mussel fauna where adequate streamside buffers are not maintained and erosion from adjacent land is allowed to enter streams (Brainwood et al. 2006, p. 511).

The range of the smooth pimpleback receives sediment from increasing levels of sedimentation from agriculture, urbanization, and sand and gravel mining (discussed in section titled Sand and Gravel Mining); sedimentation is likely to continue to threaten the smooth pimpleback.

## Dewatering

River dewatering can occur in several ways: Anthropogenic activities such as surface water diversions and groundwater pumping, and natural events, such as drought can result in mussels stranded in previously wetted areas. This is a particular concern for smooth pimpleback within and below reservoirs, where water levels are managed for various purposes that can cause water levels in the reservoir or downstream to rise or fall in very short periods of time, such as when hydropower facilities release water during peak energy demand periods. The three impoundments on the Colorado River with records of smooth pimpleback all experience periodic water level drawdowns, which may have contributed to the species' apparent extirpation from Inks Lake and Lake Marble Falls. In fact, smooth pimpleback have been found stranded (which leads to death) after drawdowns in both of these reservoirs (Howells 1996, p. 22; 1999, p. 16).

Drought can also severely impact smooth pimpleback populations. For example, the Little Brazos River, which once contained a diverse and numerous freshwater mussel community that included smooth pimpleback (Gentner and Hopkins 1966, p. 458), experienced a severe drought from about 1950 to 1956 that reduced the river to a series of small, stagnant pools. The results of this habitat degradation from the low water nearly eliminated the mussel community and killed many smooth pimpleback (Gentner and Hopkins 1966, p. 458). Later, central Texas, including the Colorado and Brazos River basins, experienced a major drought in the late 1970s (Lewis and Oliveria 1979, p. 243). Near record dry conditions in 2008 followed by a pattern of below-normal rainfall during the winter and spring of 2009 led to one of the worst droughts in recorded history for most of central Texas, including the range of the smooth pimpleback (Nielsen Gammon and McRoberts 2009, p. 2). This drought's severity was exacerbated by abnormally high air temperatures, a likely effect of climate change, which has already increased average air temperatures in Texas by at least 1°C (1.8 °F) (Nielsen-Gammon and McRoberts 2009, p. 22). Instream flows throughout the Brazos River basin during this drought were significantly reduced (USGS 2011c, p. 1) and smooth pimpleback populations in areas with reduced water levels, such as in the middle Brazos River, may have been negatively affected.

Central Texas is currently experiencing another extreme drought, with rainfall between October 2010 and July 2011 being the lowest on record during those months (LCRA 2011c, p. 1); the effects of this drought are being observed but are not yet fully known. The area suffers and continues to suffer from acute droughts, and the recent droughts of 2007-2009 and 2011 were the most severe since the all-time record drought of the 1950s (LCRA 2011c, p.1). Due to severe drought in 2011, many gravel bars that were surveyed in March and July were barely covered with water (LCRA 2011c, Photo 3, 5), large areas of the San Saba River were completely dry (LCRA 2011c, Photo 4), and the situation got even worse at the end of summer (Burlakova and Karatayev

2012b, p. 10). As the drought continued in the fall of 2011, many of the populations in San Saba River may no longer exist; 2012 was dry as well, which may have further stressed any remaining populations.

According to the National Weather Service records for 2011, more than 77 percent of Texas is experiencing moderate to extreme drought (Burlakova and Karatayev 2012b, p. 16). Current climate model simulations suggest that the American southwest could experience a 60-year stretch of heat and drought unseen since the 12th century and that the region is likely to become drier and experience more frequent droughts, with changes accelerating toward the end of the century (Woodhouse et al. 2010, pp. 21283-21288). Droughts result in a decrease in water depth and flow velocity in streams inhabited by smooth pimpleback, which reduces the availability of food and dissolved oxygen and reduces survivability. As droughts persist, mussels face hypoxia, elevated water temperature and, ultimately, death due to stranding (Golladay et al. 2004, p. 501).

### Sand and Gravel Mining

Sand and gravel mining (removing bed materials from streams) has been implicated in the destruction of mussel populations across the United States (Hartfield 1993, pp. 136–138). Sand and gravel mining causes stream instability by increasing erosion and turbidity (a measure of water clarity) and causing subsequent sediment deposition downstream (Meador and Layher 1998, pp. 8–9). These changes to the stream can result in large-scale changes to aquatic fauna, by altering habitat and affecting spawning of fish, mussels, and other aquatic species (Kanehl and Lyons 1992, pp. 4–11).

The Brazos River has a long history of sand mining, particularly in the lower river, and channel morphology changes have been attributed to destabilization due to instream sand mining in the area (USGS 2001, p. 27). The removal of sand from within the river creates sediment traps during periods of high flow, which causes scouring and erosion downstream (USGS 2001, p. 27). One gravel dredging operation in the Brazos River was documented depositing sediment as far as 1.6 km (1 mile) downstream (Forshage and Carter 1973, p. 697). Accelerated stream bank erosion and downcutting of streambeds are common effects of instream sand and gravel mining, as is the mobilization of fine sediments during sand and gravel extraction (Roell 1999, p. 7).

Within the current range of the smooth pimpleback, TPWD has issued permits for four current sand mining activities within the Brazos River (Austin, Bosque, and Fort Bend Counties) (TPWD 2004, p. 1; 2007b, p. 1, 2008b, p. 1; 2010b, p. 1). The permits allow for the repeated removal of sand and gravel at various locations within the Brazos River. The lower Brazos River, where these mining activities occur, contains one of the more numerous populations of smooth pimpleback.

In areas where repeated mining occurs, an upstream progression of channel degradation and erosion (called headcutting) can occur (Meador and Layher 1998, p. 8). Headcutting may move miles upstream in a zipper-like fashion as the upper boundary of the modified area collapses. Headcutting can be found within the majority of rivers and streams in Texas, including within the smooth pimpleback's current and historical range (Kennon et al. 1967, p. 22). Headcuts induced by sand and gravel mining can cause dramatic changes in streambank and channel shape that may



affect instream flow, water chemistry and temperature, bank stability, and siltation (Meador and Layher 1998, p. 8), all of which are harmful to freshwater mussels. Mussels are particularly vulnerable to channel degradation and sedimentation processes associated with headcutting due to their immobility (Pringle 1997, p. 429).

In addition to headcutting, mines that are located near stream channels are subject to the gravel pit being captured by the stream during flood events or due to gradual channel migration (Simmang and Curran 2006, p. 1). For example, two gravel mines along the Colorado River downstream of Austin were inundated; one by stream channel migration in 1984, one by stream capture in 1991 (Simmang and Curran 2006, p. 1). Once captured by the mainstem river, gravel mines contribute large amounts of suspended sediment to the river, causing additional turbidity and sedimentation and further degrading mussel habitat.

The smooth pimpleback population in the lower Brazos River may be currently affected by sand and gravel mining. These activities occur over a long period of time, destabilizing mussel habitat both upstream and downstream, which decreases the likelihood of recolonization after the activity has been completed. Therefore, the effects of sand and gravel mining are an ongoing threat to the smooth pimpleback and are expected to continue to occur throughout the range of the species.

### Chemical Contaminants

Chemical contaminants are ubiquitous throughout the environment and are a major reason for the decline of freshwater mussel species nationwide (Richter et al. 1997, p. 1081; Strayer et al. 2004, p. 436; Wang et al. 2007a, p. 2029). Chemicals enter the environment through both point and nonpoint discharges, including spills, industrial sources, municipal effluents, and agriculture runoff. These sources contribute organic compounds, heavy metals, pesticides, herbicides, and a wide variety of newly emerging contaminants to the aquatic environment. As a result, water quality can be degraded to the extent that mussel populations are adversely affected.

Chemical and oil spills can be especially devastating to mussels because they may result in exposure of a relatively immobile species to elevated concentrations that far exceed toxic levels. Acute and chronic exposure to oil spills in freshwater systems is largely understudied; therefore, little information is available on effects of oil spills on freshwater ecosystems (Harrel 1985, p. 223; Bhattacharyya et al. 2002, p. 205). Oil is retained much longer in marshes and other low-energy environments, such as slow-moving streams and rivers, than on wave-swept coasts (Bhattacharyya et al. 2002, p. 205). Oils have been found in sediments at low energy sites as much as 5 years after the occurrence of spills, and they may be released into the water column long after the initial spill. Oil may have various chronic effects on water-column and benthic (bottom-dwelling) species. These effects include sensory disruption, behavioral and developmental abnormalities, and reduced fertility (Bhattacharyya et al. 2002, p. 205). Oil spilled on the water surface may also limit oxygen exchange, coat the gills of aquatic organisms, and cause pathological lesions on respiratory surfaces, thereby affecting respiration in aquatic organisms. Effects of oil on freshwater mussels may result from oil settling on the sediment surfaces and accumulating in the sediment. This can prevent invertebrate colonization (Bhattacharyya et al. 2002, p. 205). Complete recovery

of benthic communities may be a matter of years, with communities in the meantime consisting solely of pollutant-tolerant organisms (Bhattacharyya et al. 2002, p. 205). Oil spills can occur from on-site accidents (tank, pipeline spills) or from tanker truck accidents within watersheds occupied by smooth pimpleback. For example, 450 gallons of oil were spilled into Lake Bastrop, a reservoir on a tributary to the Colorado River, in February 2011 (Cihock 2011, p. 1).

Examples of the exposure of smooth pimpleback to chemical contaminants include an event in 1993 when an unknown substance was dumped into a segment of the Little Brazos River upstream from a smooth pimpleback population. This site once supported an abundant and diverse number of mussel species, including the smooth pimpleback, but when it was revisited in 1993, a massive die-off of freshwater mussels had occurred (Howells 2010b, p. 11). In another instance in 2010, crude oil overflowed from a failed storage tank into Keechi Creek in Leon County, a tributary to the Navasota River (National Response Center 2010, p. 2). This location is near a small population of smooth pimpleback and upstream of one of the largest known populations of the species.

Numerous other spills have occurred within the range of the smooth pimpleback. These occurred from onsite accidents (storage tank or pipeline spills) or from tanker truck accidents within watersheds occupied by smooth pimpleback. For example, oil has spilled into the Brazos River a number of times. As much as 320,000 L (84,000 gal) of crude oil was spilled in the Brazos River in Knox County in 1991 (Associated Press 1991, p. 1). In June 2010, flooding of holding ponds adjacent to oil drilling operations leaked oil into Thompson Creek and subsequently into the Brazos River (Lewis 2010, p. 1). Also, in July 2010, oil pipelines burst and released approximately 165 barrels of crude oil into the upper Double Mountain Fork of the Brazos River in Garza County (Joiner 2010, p. 1). Although no analyses were conducted of the specific effects of these spills on smooth pimpleback, we expect that if the mussels are exposed to even moderate levels of toxic chemical contaminants, such as crude oil, adverse effects (both direct mortality and indirect effects to food source availability) are likely to occur.

Exposure of mussels to persistent low concentrations of contaminants likely to be found in aquatic environments can also adversely affect mussels and their populations. Such concentrations may not be immediately lethal, but over time can result in mortality, reduced filtration efficiency, reduced growth, decreased reproduction, changes in enzyme activity, and behavioral changes to all mussel life stages (Naimo 1995, pp. 351–352; Baun et al. 2008, p. 392). Frequently, procedures that evaluate the “safe” concentration of an environmental contaminant (for example, national water quality criteria) do not have data for freshwater mussel species or do not consider data that are available for freshwater mussels (March et al. 2007, pp. 2066–2067, 2073). One chemical that is particularly toxic to early life stages of mussels is ammonia. Sources of ammonia include agricultural activities (animal feedlots and nitrogenous fertilizers), municipal wastewater treatment plants, and industrial waste (Augspurger et al. 2007, p. 2026), as well as precipitation and natural processes (decomposition of organic nitrogen) (Goudreau et al. 1993, p. 212; Hickey and Martin 1999, p. 44; Augspurger et al. 2003, p. 2569; Newton 2003, p. 2543). Therefore, ammonia is considered a limiting factor for survival and recovery of some mussel species due to its ubiquity in aquatic environments, high level of toxicity, and because the highest concentrations typically occur in mussel microhabitats (Augspurger et al. 2003, p. 2574). In addition, studies have shown that

ammonia concentrations increase with increasing temperature and low-flow conditions (Cherry et al. 2005, p. 378; Cooper et al. 2005, p. 381), which may be exacerbated during low-flow events in streams.

In addition to ammonia, agricultural sources of chemical contaminants include two broad categories that have the potential to adversely affect mussel species: Nutrients and pesticides. High amounts of nutrients, such as nitrogen and phosphorus, in streams can stimulate excessive plant growth (algae and periphyton, among others), which in turn can reduce dissolved oxygen levels when dead plant material decomposes. Nutrient over-enrichment in streams is primarily a result of runoff of fertilizer and animal manure from livestock farms, feedlots, and heavily fertilized row crops (Peterjohn and Correll 1984, p. 1471). Over-enriched conditions are exacerbated by low-flow stream conditions, such as those experienced during typical summer season flows. Bauer (1988, p. 244) found that excessive nitrogen concentrations can be detrimental to the adult freshwater pearl mussel (*Margaritifera margaritifera*), as was evident by the positive linear relationship between mortality and nitrate concentrations. Also, a study of mussel life span and size (Bauer 1992, p. 425) showed a negative correlation between growth rate and high nutrient concentrations, and longevity was reduced as the concentration of nitrates increased. Juvenile mussels in interstitial habitats are particularly affected by depleted dissolved oxygen levels resulting from nutrient over-enrichment (Sparks and Strayer 1998, p. 133).

Mussels are also affected by metals (Keller and Zam 1991, p. 543) such as cadmium, chromium, copper, mercury, and zinc, which can negatively affect biological processes such as growth, filtration efficiency, enzyme activity, valve closure, and behavior (Keller and Zam 1991, p. 543; Naimo 1995, pp. 351–355; Jacobson et al. 1997, p. 2390; Valenti et al. 2005, p. 1244). Metals occur in industrial and wastewater effluents and are often a result of atmospheric deposition from industrial processes and incinerators. Studies have shown that copper can have toxic effects on glochidia and juvenile freshwater mussels (Wang et al. 2007a, pp. 2036–2047; Wang et al. 2007b, pp. 2048–2056). Because we know that copper contamination in water can lead to death of mussels and that these above mentioned sources are within the smooth pimpleback distribution, we conclude that the copper may be adversely affecting smooth pimpleback.

Mercury is another heavy metal that has the potential to negatively affect mussel populations, and it is widely distributed in the environment. Mercury has been detected throughout aquatic environments as a product of municipal and industrial waste and atmospheric deposition from coal burning plants. Rainbow mussel (*Villosa iris*) glochidia have been demonstrated to be more sensitive to mercury than juvenile mussels, with the median lethal concentration value of 14 parts per billion (ppb) for glochidia, compared to 114 ppb for the juvenile life stages (Valenti 2005, p. 1242). The chronic toxicity tests conducted determined that juveniles exposed to mercury greater than or equal to 8 ppb exhibited reduced growth. Acute mercury toxicity was determined to be the cause of extirpation of a diverse mussel community for a 112 km (70 mi) portion of the North Fork Holston River in Virginia (Brown et al. 2005, pp. 1455–1457). Mercury has been documented throughout Texas rivers, with particularly high concentrations in fish in the upper reaches of some of the rivers (Lee and Schultz 1994, p. 8). As with copper, we do not have information on the concentration of mercury that smooth pimpleback is being exposed to in these streams, but the

higher than expected levels in fish indicate high mercury levels in the area, which may be adversely affecting smooth pimpleback.

Pesticides are another source of contaminants in streams. Elevated concentrations of pesticides frequently occur in streams due to pesticide runoff, overspray application to row crops, and lack of adequate riparian buffers. The timing of agricultural pesticide applications in the spring often coincides with the reproductive and early life stages of mussels, which may increase the vulnerability of mussels to pesticides (Bringolf et al. 2007a, p. 2094). Little is known regarding the effect of currently used pesticides to freshwater mussels even though some pesticides, such as glyphosate (active ingredient in Roundup®), are used globally. Recent studies tested the toxicity of glyphosate, its formulations, and a surfactant (MON 0810) used in several glyphosate formulations, to early life stages of the fatmucket (*Lampsilis siliquoidea*) (Bringolf et al. 2007a, p. 2094). Studies conducted with fatmucket juveniles and glochidia determined that the surfactant was the most toxic of the compounds tested and that fatmucket glochidia were the most sensitive organisms tested to date (Bringolf et al. 2007a, p. 2094). Roundup®, technical grade glyphosate isopropylamine salt, and isopropylamine were also acutely toxic to juveniles and glochidia (Bringolf et al. 2007a, p. 2097). These commonly applied pesticides may be adversely affecting smooth pimpleback populations.

The effects of other widely used pesticides, including atrazine, chlorpyrifos, and permethrin, on glochidia and juvenile life stages have also recently been studied (Bringolf et al. 2007b, p. 2101). Environmentally relevant concentrations (concentrations that may be found in streams) of permethrin and chlorpyrifos were found to be toxic to glochidia and juvenile fatmucket (Bringolf et al. 2007b, pp. 2104–2106). Commonly applied pesticides are a threat to mussels as a result of their widespread use. All of these pesticides are commonly used on agricultural lands throughout the range of the smooth pimpleback, which may be adversely affecting the species.

A potential, but undocumented, threat to freshwater mussels, including smooth pimpleback, are compounds referred to as “emerging contaminants” that are being detected in aquatic ecosystems at an increasing rate. These include pharmaceuticals, hormones, and other organic contaminants that have been detected downstream from urban areas and livestock production (Kolpin et al. 2002, p. 1202) and have been shown to affect fish behavior (TCEQ 2010b, p. 3). In samples of the Trinity River, for example, compounds such as antidepressants, antihistamines, blood pressure lowering medication, antiseizure medication, and antimicrobial compounds were all detected during a 2006 study (TCEQ 2010b, pp. 27–28). A large potential source of these emerging contaminants is wastewater being discharged through both permitted (National Pollutant Discharge Elimination System (NPDES)) and nonpermitted sites within the Colorado and Brazos River systems. Although streams within the range of smooth pimpleback have not been tested for these emerging contaminants, permitted discharge sites are ubiquitous in watersheds with smooth pimpleback populations, providing many opportunities for contaminants to impact the species.

A study in the Blanco River found that mussels may be adversely affected by sewage effluent (Horne and McIntosh 1979, p. 132). Ammonia levels below the outfall were three times higher than the levels above the outfall and were higher than recently determined toxicity values of ammonia

for mussels (Augsperger et al. 2003, p. 2572). The river was nutrient-enriched for miles downstream, and mussels were less abundant below the outfall than above (Horne and McIntosh 1979, pp. 124– 125, 132).

Texas Commission on Environmental Quality (TCEQ) data for 2010 indicated that 26 of the 98 assessed water bodies within Colorado River basin and 81 of approximately 124 assessed water bodies within Brazos River basin did not meet surface water quality standards and were classified as impaired water bodies (Texas Clean Rivers Program 2010a, p. 5; TCEQ 2010c, pp. 1–106). These water bodies were impaired with dissolved solids, nitrites, nitrates, bacteria, low dissolved oxygen, aluminum, sulfates, selenium, chloride, orthophosphorus, phosphorus, Chlorophyll a, and low pH associated with agricultural, urban, municipal, and industrial runoff. Of these, nitrites and low dissolved oxygen are known to be harmful to freshwater mussels. Agricultural pesticides and emerging contaminants are likely also present in streams inhabited by smooth pimpleback. There are 53 wastewater treatment plants permitted to discharge more than one million gallons per day into the Brazos River basin (Valenti and Brooks 2008, p. 12); the outfalls of these treatment plants have not been tested to determine if they contain contaminants of note.

Releases of chemical contaminants, such as oil, ammonia, copper, mercury, nutrients, pesticides, and other compounds into the habitat of the smooth pimpleback are an ongoing threat to the smooth pimpleback. The species is vulnerable to acute contamination from spills, as well as chronic contaminant exposure, which has occurred and is expected to continue to occur throughout the range of the smooth pimpleback.

#### Summary of Factor A

The reduction in numbers and range of the smooth pimpleback is primarily the result of the long-lasting effects of habitat alterations such as the effects of impoundments, sedimentation, dewatering, sand and gravel mining, and chemical contaminants. Impoundments occur throughout the range of the species and have farreaching effects to riverine habitat both upstream and downstream of the dams. Both the Colorado and Brazos River systems have experienced a large amount of sedimentation from agriculture, instream mining, and urban development. Sand and gravel mining affects smooth pimpleback habitat by increasing sedimentation and channel instability downstream and by causing headcutting upstream. Chemical contaminants exceeding the standards developed to support aquatic life have been documented throughout the range of the species and may represent a significant threat to the smooth pimpleback. However, the large populations in the San Saba River, lower Brazos River, Navasota River, Leon River, and Yegua Creek indicate that some smooth pimpleback populations are not currently as vulnerable to habitat loss as others. Therefore, based upon our review of the best commercial and scientific data available, we conclude that the present or threatened destruction, modification, or curtailment of its habitat or range is an immediate threat of moderate magnitude to the smooth pimpleback.

#### **B. Overutilization for commercial, recreational, scientific, or educational purposes:**

The smooth pimpleback is not a commercially valuable species and has never been harvested in Texas as a commercial mussel species (Howells 2010b, p.12). Some scientific collecting occurs but is not likely to be a significant threat to the species because it occurs only rarely. However, handling mussels can disturb gravid females and result in glochidial loss and subsequent reproductive failure. Additionally, handling has also been shown to reduce shell growth across mussel species, including several species of *Lampsilis* (Haag and Commens-Carson 2008, pp. 505–506). Repeated handling by researchers may adversely affect smooth pimpleback individuals, but these activities are occurring rarely and are not likely to be a threat to populations. Handling for scientific purposes contributes to the long-term conservation of the species. We do not have any evidence of risks to the smooth pimpleback from overutilization for commercial, recreational, scientific, or educational purposes, and we have no reason to believe this factor will become a threat to the species in the future. Based upon the best scientific and commercial information available, we conclude that overutilization for commercial, recreational, scientific, or educational purposes does not pose a threat to the smooth pimpleback rangewide.

### **C. Disease or predation:**

#### **Disease**

Little is known about disease in freshwater mussels. However, disease is believed to be a contributing factor in documented mussel die-offs in other parts of the United States (Neves 1987, pp. 11–12). Diseases have not been documented or observed during any studies of smooth pimpleback.

#### **Predation**

Raccoons will prey on freshwater mussels stranded by low waters or deposited in shallow water or on bars following flooding or low water periods (Howells 2010c, p. 12). Predation of smooth pimpleback by raccoons may be occurring occasionally, but there is no indication it is a significant threat to the status of the species. Some species of fish feed on mussels, such as common carp, freshwater drum, and redear sunfish, all of which are common throughout the range of smooth pimpleback (Hubbs et al. 2008, pp. 19, 45, 53). Common species of flatworms are voracious predators of newly metamorphosed juvenile mussels of many species (Zimmerman et al. 2003, p. 30). Predation is a normal factor influencing the population dynamics of a healthy mussel population; however, predation may amplify declines in small populations primarily caused by other factors.

#### **Summary of Factor C**

Disease in freshwater mussels is poorly known, and we do not have any information indicating it is a threat to the smooth pimpleback. Additionally, predation is a natural ecological interaction and we have no information indicating the extent of any predation is a threat to populations of smooth pimpleback. Based upon the best scientific and commercial information available, we conclude that disease or predation is not a threat to the smooth pimpleback.

## **D. The inadequacy of existing regulatory mechanisms:**

The Act requires us to examine the adequacy of existing regulatory mechanisms with respect to threats that may place the smooth pimpleback in danger of extinction or increase its likelihood of becoming so in the future. Existing regulatory mechanisms that could affect threats to the smooth pimpleback include State and Federal laws such as the Texas Threatened and Endangered Species regulations, Texas freshwater mussel sanctuaries, State and Federal sand and gravel mining regulations, and regulation of point and non-point source pollution.

### **Texas Threatened and Endangered Species Regulations**

On January 8, 2010, the Texas Parks and Wildlife Commission placed 15 species of freshwater mussels, including the smooth pimpleback, on the State threatened list (Texas Register 2010, pp. 6–10). Section 68.002 of the Texas Parks and Wildlife (TPW) Code and Section 65.171 of the Texas Administrative Code (TAC) prohibit the direct take of a threatened species, except under issuance of a scientific collecting permit. “Take” is defined in Section 1.101(5) of the TPW Code as collect, hook, hunt, net, shoot, or snare, by any means or device, and includes an attempt to take or to pursue in order to take. While this law protects individuals from take, it is difficult to enforce and does not provide any protection for smooth pimpleback habitat. Moreover, our assessment finds that the species is not threatened by take (see Factor B above). There are no State provisions under the Texas Threatened and Endangered Species Regulations for reducing or eliminating the threats (see Factor A above) that may adversely affect smooth pimpleback or its habitat. In addition, these State regulations do not call for development of a recovery plan that will restore and protect existing habitat for the species. For these reasons, we find that existing Texas regulatory mechanisms for State-listed threatened species are currently inadequate to protect smooth pimpleback and its habitat or to prevent further decline of the species.

### **Freshwater Mussel Sanctuaries**

The TPWD has designated specific areas of streams and reservoirs as no harvest mussel sanctuaries (31 TAC, part 2, chapter 57, subpart B, Rule 57.157). The locations of the designated mussel sanctuaries were selected because they support populations of rare and endemic mussel species or are important for maintaining, repopulating, or allowing recovery of mussels in watersheds where they have been depleted. Unfortunately, mussel sanctuaries only restrict the harvest of mussels and do not address other activities that may affect mussels or their habitats. Therefore, these designations provide no regulatory mechanisms to protect smooth pimpleback from habitat alteration.

### **State Sand and Gravel Mining Regulations**

The TPWD has been responsible for regulating the “disturbance of taking” streambed materials since 1911 (Meador and Layher 1998, p. 11) and has issued several permits for ongoing activities within the smooth pimpleback range (for more information on the effects of sand and gravel mining

on smooth pimpleback, please refer to “Sand and Gravel Mining” under Factor A in Five-Factor Evaluation). Sand and gravel mining may be one of the least regulated of all mining activities (Meador and Layher 1998, p. 10).

## Clean Water Act

The U.S. Army Corps of Engineers (USACE) retains oversight authority and requires a permit for gravel and sand mining activities that deposit fill into streams under section 404 of the Clean Water Act (33 U.S.C. 1251 et seq.). Additionally, a permit is required under section 10 of the Rivers and Harbors Act (33 U.S.C. 401 et seq.) for navigable waterways. However, many mining operations do not fall under these two categories. For example, nationwide permits are issued by the USACE for types of projects that are presumed to have minimal environmental impacts. However, projects permitted by nationwide permits, such as small mining operations, may have cumulative effects on aquatic species like the smooth pimpleback through increased sedimentation and channel instability.

Point source discharges of potential contaminants within the range of the smooth pimpleback have been reduced since the inception of the Clean Water Act, but this reduction may not provide adequate protection for filter-feeding organisms that can be affected by extremely low levels of contaminants (see “Chemical Contaminants” under Factor A). The EPA’s established water quality criteria may not be protective of mussels. Current water quality standards applied by EPA were established to be protective of aquatic life; however, freshwater mussels were not used to develop these standards (EPA 2005, p. 5), and current research reveals mussels to be more sensitive to many aquatic pollutants than the tested organisms (Augsperger et al. 2007, p. 2025). For example, Augspurger et al. (2003, p. 2572) and Sharpe (2005, p. 28) suggested that the criteria for ammonia may not be sufficient to prevent impacts to mussels under current and future climate conditions. In addition, chronic copper concentrations lethal to juvenile freshwater mussels have been shown to be less than the EPA’s 1996 chronic water quality criterion for copper (Wang et al. 2007b, pp. 2052–2055). Based on this information, the existing EPA water quality criteria may not be sufficient to prevent negative effects to the smooth pimpleback.

Nonpoint source pollution such as sedimentation and chemical contamination is considered a significant threat to smooth pimpleback habitat; however, the Clean Water Act does not adequately protect smooth pimpleback habitat from nonpoint source pollution, because most activities that cause nonpoint source pollution are not regulated under the Clean Water Act.

## Summary of Factor D

Despite some State and Federal laws protecting the species and water quality, the smooth pimpleback continues to decline due to the effects of habitat destruction, poor water quality, contaminants, and other factors. The regulatory measures described above are not sufficient to significantly reduce or remove the threats to the smooth pimpleback. Based upon our review of the best commercial and scientific data available, we conclude that the lack of existing regulatory mechanisms is an immediate threat of moderate magnitude to the smooth pimpleback.



## **E. Other natural or manmade factors affecting its continued existence:**

Natural and manmade factors that threaten the smooth pimpleback include climate change, population fragmentation and isolation, and nonnative species.

### **Climate Change**

It is widely accepted that changes in climate are occurring worldwide (International Panel on Climate Change (IPCC) 2007, p. 30). Understanding the effects of climate change on the smooth pimpleback is important because the disjunct nature of the remaining smooth pimpleback populations, coupled with the limited ability of mussels to migrate, makes it unlikely that the smooth pimpleback can adjust its range in response to changes in climate (Strayer 2008, p. 30). For example, changes in temperature and precipitation can increase the likelihood of flooding or increase drought duration and intensity, resulting in direct effects to freshwater mussels like the smooth pimpleback (Hastie et al. 2003, pp. 40–43; Golloday et al. 2004, p. 503). Because the range of the smooth pimpleback has been reduced to isolated locations with low population numbers in small to medium sized rivers and streams, the smooth pimpleback is vulnerable to climatic changes that could decrease the availability of water or produce more frequent scouring flood events. Indirect effects of climate change may include declines in host fish populations, habitat reduction, and changes in human activity in response to climate change (Hastie et al. 2003, pp. 43–44).

For the next two decades, a warming of about 0.2 °C (0.4 °F) per decade is projected across the United States (IPCC 2007, p. 12), and hot extremes, heat waves, and heavy precipitation and flooding are expected to increase in frequency (IPCC 2007, p. 18). As with many areas of North America, central Texas is projected to experience an overall warming trend in the range of 2.5 to 3.3 °C (4.5 to 6 °F) over the next 50 to 200 years (Mace and Wade 2008, p. 656). Even under lower greenhouse gas emission scenarios, recent projections forecast a 2.8 °C (5 °F) increase in temperature and a 10 percent decline in precipitation in central Texas by 2080 to 2099 (Karl et al. 2009, pp. 123–124). Based on our current understanding of climate change, air temperatures are expected to rise and precipitation patterns are expected to change in areas occupied by the smooth pimpleback. Karl et al. (2009, p. 12) also suggests that climate change impacts on water resources in the southern Great Plains (including central Texas) are expected as rising temperatures and decreasing precipitation exacerbate an area already plagued by low rainfall, high temperatures, and unsustainable water use practices.

One preliminary study forecasting the possible hydrological impacts of climate change on the annual runoff and its seasonality in the upper Colorado River watershed was conducted by CH2M HILL (2008). In this initial evaluation, four modeling scenarios (chosen to represent a range of possible future climatic conditions) were each run under a 2050 and 2080 time scenario, producing annual surface water runoff estimates at multiple sites with stream gages in the Colorado River basin. For the 2050 scenarios, the results from all four climate change scenarios predicted

significant decreases in annual runoff totals compared to historic averages (CH2M HILL 2008, pp. 7–30—7–32). For the 2080 scenarios, one model predicted increases in annual runoff; the other three 2080 scenarios predicted decreases in annual runoff (CH2M HILL 2008, pp. 7–30—7–33). The modeling efforts from this study focus on annual averages and cannot necessarily account for the seasonal variations in flooding events or long periods of drought. However, the study demonstrates the potential effects of climate change on surface water availability, which is forecasted to result in an overall decline in stream flows in the region where the smooth pimpleback occurs.

In summary, climate change could affect the smooth pimpleback through the combined effects of global and regional climate change, along with the increased probability of long-term drought. Climate change exacerbates threats such as habitat degradation from prolonged periods of drought, increased water temperature, and the increased allocation of water for municipal, agricultural, and industrial use. As such, climate change, in and of itself, may affect the smooth pimpleback, but the magnitude and imminence (when the effects occur) of the effects remain uncertain. Based upon our review of the best commercial and scientific data available, we conclude that the effects of climate change in the future will likely exacerbate the current and ongoing threats of habitat loss and degradation caused by other factors, as discussed above.

#### Population Fragmentation and Isolation

As with many freshwater mussels, several of the remaining populations of the smooth pimpleback are small and geographically isolated and thus are susceptible to genetic drift, inbreeding depression, and random or chance changes to the environment, such as toxic chemical spills (Watters and Dunn 1995, pp. 257–258), or dewatering. Inbreeding depression can result in death, decreased fertility, smaller body size, loss of vigor, reduced fitness, and various chromosomal abnormalities (Smith 1974, pp. 350). Despite any evolutionary adaptations for rarity, habitat loss and degradation increase a species' vulnerability to extinction (Noss and Cooperrider 1994, pp. 58–62). Numerous authors (including Noss and Cooperrider 1994, pp. 58–62; Thomas 1994, p. 373) have indicated that the probability of extinction increases with decreasing habitat availability. Although changes in the environment may cause populations to fluctuate naturally, small and low-density populations are more likely to fluctuate below a minimum viable population (the minimum or threshold number of individuals needed in a population to persist in a viable state for a given interval) (Gilpin and Soule 1986, pp. 25–33; Shaffer 1981, p. 131; Shaffer and Samson 1985, pp. 148–150).

Historically, the smooth pimpleback was widespread throughout much of the Colorado and Brazos River systems when few natural barriers existed to prevent migration (via host species) among suitable habitats. Construction of dams, however, likely destroyed many smooth pimpleback populations through drastic habitat changes and isolated the remnant populations from each other. The extensive impoundment of the Brazos and Colorado River basins has fragmented smooth pimpleback populations throughout these river systems. For fertilization, smooth pimpleback females need an upstream male to release sperm; populations with few individuals reduce the likelihood that females will be exposed to sperm while siphoning. Therefore, recruitment failure is a

potential problem for many small populations rangewide, a potential condition exacerbated by its reduced range and increasingly isolated populations. If downward population trends continue, further significant declines in total smooth pimpleback population size and consequent reduction in long-term survivability may soon become apparent. Small smooth pimpleback populations, including those in Lake LBJ Reservoir and the middle Brazos, Little, and Little Brazos Rivers, may be below the minimum population size required to maintain population viability into the future, therefore making these populations more vulnerable to extirpation since they are less likely to be able to recover through recruitment from events that reduce but do not extirpate populations.

Additionally, these small populations are more vulnerable to extirpation from stochastic (random) natural events, as the lack of connectivity among populations does not permit nearby populations to recolonize areas affected by intense droughts, toxic spills, or other isolated events that result in significant mussel die-offs. When species are limited to small, isolated habitats, as the smooth pimpleback is, they are more likely to become extinct due to a local event that negatively effects the population (McKinney 1997, p. 497; Minckley and Unmack 2000, pp. 52–53; Shepard 1993, pp. 354–357). While the populations' small, isolated nature does not represent an independent threat to the species, it does substantially increase the risk of extirpation from the effects of all other threats, including those addressed in this analysis, and those that could occur in the future from unknown sources.

Based upon our review of the best commercial and scientific data available, we conclude that fragmentation and isolation of small remaining populations of the smooth pimpleback are occurring and are ongoing threats to the species throughout all of its range. Further, stochastic events may play a magnified role in extirpation of small, isolated populations.

### Nonnative Species

Various nonnative species of aquatic organisms are firmly established within the range of the smooth pimpleback and pose a threat to the species. Golden algae (*Prymnesium parvum*) is a microscopic algae considered to be one of the most harmful algal species to fish and other gill-breathing organisms (Lutz-Carrillo et al. 2010, p. 24). Golden algae was first discovered in Texas in 1985 and is presumed to have been introduced from western Europe (Lutz-Carrillo et al. 2010, p. 30). Since its introduction, golden algae have been found in Texas rivers and lakes, including two lakes in central Texas (Baylor University 2009, p. 1). Under certain environmental conditions, golden algae can produce toxins that can cause massive fish and mussel kills (Barkoh and Fries 2010, p. 1; Lutz-Carrillo et al. 2010, p. 24). Evidence shows that golden algae probably caused fish kills in Texas as early as the 1960s, but the first documented fish kill due to golden algae in inland waters of Texas occurred in 1985 on the Pecos River in the Rio Grande basin (TPWD 2002, p. 1). The range of golden algae has increased to include portions of the Brazos and Colorado River basins, among others, and it has been responsible for killing more than 8 million fish in the Brazos River since 1981 and more than 2 million fish in the Colorado River since 1989 (TPWD 2010a, p. 1). Although actual mussel kills in Texas due to golden algae have not been

recorded in the past, the toxin can kill mussels. Therefore, the elimination of host fish and the poisonous nature of the toxin to mussels make future golden algae blooms a threat to the smooth pimpleback.

An additional nonnative species, the zebra mussel (*Dreissena polymorpha*) are a potential threat to the smooth pimpleback. This invasive species has been responsible for the extirpation of freshwater mussels in other regions of the United States, including the Higgin's eye (*Lampsilis higginsii*) in Wisconsin and Iowa (Service 2006, pp. 9–10). Zebra mussels attach in large numbers to the shells of live native mussels and are implicated in the loss of entire native mussel beds (Ricciardi et al. 1998, p. 615). This fouling impedes locomotion (both laterally and vertically), interferes with normal valve movements, deforms valve margins, and essentially suffocates and starves the native mussels by depleting the surrounding water of oxygen and food (Strayer 1999, pp. 77–80). Heavy infestations of zebra mussels on native mussels may overly stress the animals by reducing their energy reserves. Zebra mussels may also filter the sperm and possibly glochidia of native mussels from the water column, thus reducing reproductive potential. Habitat for native mussels may also be degraded by large deposits of zebra mussel pseudofeces (undigested waste material passed out of the incurrent siphon) (Vaughan 1997, p. 11).

Zebra mussels are currently found within the range of the smooth pimpleback. A live adult zebra mussel was first documented in Lake Texoma on the Red River (on the north Texas border with Oklahoma) in 2009 (TPWD 2009a, p. 1). Since that time, additional zebra mussels have been reported from Lake Texoma, where they are now believed to be well established (TPWD 2009c, p. 1). In spring and summer of 2013, Texas Parks and Wildlife Department (TPWD) monitored 23 other Texas reservoirs and found that zebra mussels may be present in two additional reservoirs: Lake Worth and Joe Pool. No adult zebra mussels or veligers have been found in either of the aforementioned water bodies (TPWD 2013, p. 1). To date, Lake Texoma, Lake Ray Roberts, Lake Lewisville, Lake Bridgeport, Lake Lavon, Lake Belton, and Lake Waco reservoirs and Elm Fork of the Trinity River are known to harbor zebra mussels (TPWD 2014, p. 1). Zebra mussels are likely to spread to many other Texas reservoirs through accidental human transport (Schneider et al. 1998, p. 789). Although zebra mussels tend to proliferate in reservoirs or large pools, released zebra mussel veligers float downstream and attach to any hard surface available, rendering downstream smooth pimpleback populations extremely vulnerable to attachment and fouling. Because zebra mussels are so easily introduced to new locations, the potential for zebra mussels to continue to expand in Texas and further invade the range of the smooth pimpleback is high. If this occurs, the smooth pimpleback is vulnerable to zebra mussel attachment and subsequent deprivation of oxygen, food, and mobility.

A molluscivore (mollusk eater), the black carp (*Mylopharyngodon piceus*) is a potential threat to the smooth pimpleback. The species has been commonly used by aquaculturists to control snails or for research in fish production in several States, including Texas (72 FR 59019, October 18, 2007). Black carp can reach more than 1.3 m (4 ft) in length and 150 pounds (68 kilograms (kg)) (Nico and Williams 1996, p. 6). Foraging rates for a 4-year-old fish average 3 to 4 pounds (1.4 to 1.8 kg) a day, indicating that a single individual could consume 10 tons (9,072 kg) of native mollusks over its lifetime (Mississippi Interstate Cooperative Resource Association (MICRA) 2005, p. 1). Black carp

can escape from aquaculture facilities. For example, in 1994 30 black carp escaped from an aquaculture facility in Missouri during a flood. Other escapes into the wild by non-sterile carp are likely to occur. Because of the high risk to freshwater mussels and other native mollusks, the Service recently listed black carp as an injurious species under the Lacey Act (72 FR 59019, October 18, 2007), which prevents importations and interstate transfer of this harmful species, but does not prevent its release into the wild once it is in the State. If the black carp were to escape within the range of the smooth pimpleback, it would likely negatively affect native mussels, including the smooth pimpleback.

Based upon our review of the best commercial and scientific data available, we conclude that golden algae is an ongoing threat to the smooth pimpleback, and other nonnative species, such as zebra mussels and black carp, are a potential future threat to the smooth pimpleback that is likely to increase as these exotic species expand their occupancy within the range of the smooth pimpleback.

#### Summary of Factor E

The effects of climate change, while difficult to quantify at this time, are likely to exacerbate the current and ongoing threat of habitat loss caused by other factors, and the small sizes and fragmented nature of the remaining populations render them more vulnerable to extirpation. In addition, nonnative species, such as golden algae, currently threaten the smooth pimpleback, and the potential introduction of zebra mussels and black carp are potential future threats. Based upon our review of the best commercial and scientific data available, we conclude that other natural or manmade factors are immediate threats of moderate magnitude to the smooth pimpleback.

### **Conservation Measures Planned or Implemented :**

The smooth pimpleback is listed as threatened by the TPWD in Texas and is a high priority species in the Texas Wildlife Action Plan 2005-2010 (TPWD 2005, p. 756). The Service, TPWD, academia, and other resource agencies have proposed and ongoing studies in Texas' river systems for Texas freshwater mussels, including the smooth pimpleback, observing life history parameters (including determination of ecological fish hosts), survivability of juveniles, monitoring habitat, and analyzing population dynamics. In addition, TPWD has established a Mussel Watch group.

The Service is currently working on forming and implementing the use of a Strategic Conservation Plan for Texas Freshwater Mussels that will result in additional conservation measures such as, best management practices, survey protocols, relocation protocols, and monitoring guidelines. The Service will be collaborating with other Federal, State, and non-governmental agencies during the formation and implementation of the Strategic Conservation Plan.

### **Summary of Threats :**

This assessment identifies threats to the smooth pimpleback attributable to Factors A, D, and E. The primary threat to the species is from habitat destruction and modification (Factor A) from

impoundments, which scour riverbeds, thereby removing mussel habitat, decreases water quality, modifies stream flows, and restricts fish host migration and distribution of freshwater mussels. Additional threats under Factor A include sedimentation, dewatering, sand and gravel mining, and chemical contaminants. Also, most of these threats may be exacerbated by the current and projected effects of climate change (discussed under Factor E). Threats to the smooth pimpleback are not being adequately addressed through existing regulatory mechanisms (Factor D). Because of the limited distribution of this endemic species and its lack of mobility, these threats are likely to lead to the extinction of the smooth pimpleback in the foreseeable future.

**For species that are being removed from candidate status:**

\_\_\_\_\_ Is the removal based in whole or in part on one or more individual conservation efforts that you determined met the standards in the Policy for Evaluation of Conservation Efforts When Making Listing Decisions(PECE)?

**Recommended Conservation Measures :**

Continued survey and monitoring efforts are needed throughout former and occupied sites to better define the species' distribution and status in the Colorado and Brazos River systems.

Continued biological and ecological research efforts are needed to identify host fish, spawning and brooding seasons, glochidia, and habitat and physiochemical parameters for the smooth pimpleback.

The Service will continue to work with TPWD, USGS, and others needed research in order to facilitate the conservation and preservation of the smooth pimpleback.

Long-term conservation measures need to be developed to facilitate and accomplish cooperative efforts between resource management agencies and private landowners. The development candidate conservation agreements (with assurances) with interested parties would initiate conservation for the smooth pimpleback.

The Service will continue working with resource management agencies and the Texas Department of Transportation (TxDOT) on developing best management practices for proposed adjacent and instream impacts specific to Texas water systems.

The Service will continue working with resource management agencies and academia on developing a drought contingency plan that will facilitate the management and monitoring of mussel populations that harbor species of concern (i.e. the smooth pimpleback) during times of drought.

The Service will continue working with resource management agencies, TxDOT, and academia on the development of standard mussel survey, relocation, and monitoring protocols, which would establish a commonality among the wide variety of methods currently being used in Texas and would establish a baseline of what kind of data needs to be collected while conducting surveys.

## Priority Table

Magnitude	Immediacy	Taxonomy	Priority
High	Imminent	Monotypic genus	1
		Species	2
		Subspecies/Population	3
	Non-imminent	Monotypic genus	4
		Species	5
		Subspecies/Population	6
<b>Moderate to Low</b>	<b>Imminent</b>	Monotypic genus	7
		<b>Species</b>	<b>8</b>
		Subspecies/Population	9
	Non-Imminent	Monotype genus	10
		Species	11
		Subspecies/Population	12

### Rationale for Change in Listing Priority Number:

No change in listing priority number.

### Magnitude:

We consider the threats that the smooth pimpleback faces to be moderate in magnitude. Habitat loss and degradation from impoundments, sedimentation, sand and gravel mining, and chemical contaminants are widespread throughout the range of the smooth pimpleback, but several large populations remain, including one that was recently discovered, indicating the threats are not high in magnitude.

### Imminence :

We consider the threats to the smooth pimpleback as described under Factor A D and E under the Five-Factor Evaluation for Smooth Pimpleback to be imminent because these threats are ongoing and will continue in the foreseeable future. Habitat loss and destruction has already occurred and will continue as the human population continues to grow in central Texas. Several smooth pimpleback populations may already be below the minimum viable population requirement, which

would cause a reduction in the number of populations and an increase in the species' vulnerability to extinction. These threats are exacerbated by climate change, which will increase the frequency and magnitude of droughts. Therefore, we consider these threats to be imminent.

☐ Yes Have you promptly reviewed all of the information received regarding the species for the purpose of determination whether emergency listing is needed?

## **Emergency Listing Review**

☐ No Is Emergency Listing Warranted?

## **Description of Monitoring:**

The Texas Parks and Wildlife Department's Mussel Watch group has been conducting surveys throughout Texas and found several fresh dead smooth pimpleback in the Colorado and Brazos River systems. The groups continued efforts along with historic data has sparked the interest of academia to further survey efforts in the Colorado and Brazos River systems where a couple of large, stable, reproducing populations were discovered and are now being closely monitored. These recent discoveries will likely lead to increased survey and monitoring efforts throughout Texas.

**Indicate which State(s) (within the range of the species) provided information or comments on the species or latest species assessment:**

none

**Indicate which State(s) did not provide any information or comment:**

Texas

## **State Coordination:**

## **Literature Cited:**

Allan, J. D. and A. S. Flecker. 1993. Biodiversity conservation in running waters. BioScience 43:32-43.

Arey, L. B. 1932. The formation and structure of the glochidial cyst. Biological Bulletin 62:212-221.

Armour, C., D. Duff, and W. Elmore. 1994. The effects of livestock grazing on western riparian and stream ecosystem. Fisheries 19(9):9-12.

Associated Press. 1991. 84,000 gallons of crude oil spill into Brazos River. Houston Chronicle June 9, 1991.



[http://www.chron.com/CDA/archives/archive.mpl/1991\\_788384/84-000-gallons-of-crude-oil-spill-into-l](http://www.chron.com/CDA/archives/archive.mpl/1991_788384/84-000-gallons-of-crude-oil-spill-into-l)  
Accessed July 12, 2011.

Athearn, H. 1970. Discussion of Dr. Heard's paper. American Malacological Union Symposium on Rare and Endangered Mollusks. *Malacologia* 10:28-31.

Augspurger, T., F. J. Dwyer, C. G. Ingersoll, and C. M. Kane. 2007. Advances and opportunities in assessing contaminant sensitivity of freshwater mussel (Unionidae) early life stages. *Environmental Toxicology and Chemistry* 26:2025-2028.

Augspurger, T., A. E. Keller, M. C. Black, W. G. Cope, and F. J. Dwyer. 2003. Water quality guidance for protection of freshwater mussels (Unionidae) from ammonia exposure. *Environmental Toxicology and Chemistry* 22:2569-2575.

Austin City Connection. 2011. City of Austin demographics.  
<http://www.ci.austin.tx.us/demographics/> Accessed August 19, 2011.

Baird, M. S. 2000. Life history of the spectaclecase, *Cumberlandia monodonta* Say, 1829 (Bivalvia, Unionoidea, Margaritiferidae). Unpublished master's thesis, Missouri State University, Springfield. 108 pp.

Barkoh, A. and L. T. Fries. 2010. Aspects of the origins, ecology, and control of golden alga *Prymnesium parvum*: introduction to the featured collection. *Journal of the American Water Resources Association* 46:1-5.

Bauer, G. 1988. Threats to the freshwater pearl mussel *Margaritifera margaritifera* L. in central Europe. *Biological Conservation* 45:239-254.

Bauer, G. 1992. Variation in the life span and size of the freshwater pearl mussel. *Journal of Animal Ecology* 61:425-436.

Baun, A., N. B. Hartmann, K. Grieger, and K. O. Kusk. 2008. Ecotoxicity of engineered nanoparticles to aquatic invertebrates: a brief review and recommendations for future toxicity testing. *Ecotoxicology* 17:387-395.

Baylor University. 2009. Baylor researchers identify what makes deadly algae more toxic. Available at: <http://www.baylor.edu/pr/news.php?action=story&story=64323> Accessed June 22, 2011.

Bhattacharyya, S., P. L. Klerks, and J. A. Nyman. 2003. Toxicity to freshwater organisms from oils and oil spill chemical treatments in laboratory microcosms. *Environmental Pollution* 122:205-215.

Bogan, A. E. 1993. Freshwater bivalve extinctions (Mollusca: Unionoida): a search for causes. *American Zoologist* 33:599-609.

Bogan, A. E. 2011. Phone conversation regarding resurrected genus *Amphinaias*. North Carolina Museum of Natural History, Raleigh. June 10, 2011.

Brainwood, M., S. Burgin, and M. Byrne. 2006. Is the decline of freshwater mussel populations in a regulated coastal river in south-eastern Australia linked with human modification of habitat? *Aquatic Conservation: Marine and Freshwater Ecosystems* 16:501-516.

Bramlette, D. and P. Cosel. 2010. Austin cleaning up big wastewater spill. KXAN News, August 31, 2010. <http://www.kxan.com/dpp/elections/local/wastewater-spill-discovered> Accessed July 12, 2011

Brazos G Regional Water Planning Group. 2010. 2011 Brazos G regional water plan. Administered by the Brazos River Authority, prepared by HDR Engineering, Inc. September 2010. Available at: [http://www.twdb.state.tx.us/wrpi/rwp/3rdRound/2011\\_RWP/RegionG/](http://www.twdb.state.tx.us/wrpi/rwp/3rdRound/2011_RWP/RegionG/).

Brazos River Authority. 2006. Targeted total suspended solids stormwater sampling in the Brazos River watershed downstream of Lake Possum Kingdom. Special Studies Final Report, December 15, 2006. 12 pp.

Brazos River Authority. 2007. Basin overview. 6 pp. Available at: [www.brazos.org/crpPDF/BasinOverview\\_2007.pdf](http://www.brazos.org/crpPDF/BasinOverview_2007.pdf) Accessed June 16, 2011.

Brim Box, J., and J. Mossa. 1999. Sediment, land use, and freshwater mussels: prospects and problems. *Journal of the North American Benthological Society* 18:99-117.

Bringolf, R. B., W. G. Cope, S. Mosher, M. C. Barnhart, and D. Shea. 2007a. Acute and chronic toxicity of glyphosate compounds to glochidia and juveniles of *Lampsilis silquoidea* (Unionidae). *Environmental Toxicology and Chemistry* 26:2094-2100.

Bringolf, R. B., W. G. Cope, M. C. Barnhart, S. Mosher, P. R. Lazaro, and D. Shea. 2007b. Acute and chronic toxicity of pesticide formulations (atrazine, chlorpyrifos, and permethrin) to glochidia and juveniles of *Lampsilis silquoidea*. *Environmental Toxicology and Chemistry* 26:2101-2107.

Brown, M. E., M. Kowalski, R. J. Neves, D. S. Cherry, and M. E. Schreiber. 2005. Freshwater mussel shells as environmental chronicles: geochemical and taphonomic signatures of mercury-related extirpations in the North Fork Holston River, Virginia. *Environmental Science and Technology* 39:1455-1462.

Burlakova, L.E. and A.Y. Karatayev. 2008. Performance report—Interim: State-Wide Assessment of Unionid Diversity in Texas. Texas State Wildlife Grants Program, Federal Aid Grant No. T-43, 13 pp.

Burlakova, L. E. and A. Y. Karatayev. 2009. Performance report – interim: state-wide assessment of unionid diversity in Texas. Texas State Wildlife Grants Program, Federal Aid Grant No. T-43. 8 pp.

Burlakova, L. E. and A. Y. Karatayev. 2010a. Performance report – final: state-wide assessment of unionid diversity in Texas. Texas State Wildlife Grants Program, Federal Aid Grant No. T-43. 30 pp.

Burlakova, L. E. and A. Y. Karatayev. 2010b. Database of rare mussel collections in Texas, 2005 – 2008. Texas Parks and Wildlife Department.

Burlakova, L. E. and A. Y. Karatayev. 2011. Update on the status of rare and endemic species in Texas undergoing full 12-month status reviews (March 2011). Preliminary report of survey of threatened freshwater mussels (Bivalvia: Unionidae) in Texas. Texas State Wildlife Grants Program. 8 pp.

Burlakova, L.E. and A.Y. Karatayev. 2012a. Update on the Status of rare and endemic Species in Texas for IUCN. 23 pp.

Burlakova, L.E. and A.Y. Karatayev. 2012b. Performance Report—Final: state-wide assessment of unionid diversity in Texas. Texas State Wildlife Grants Program, Federal Aid Grant No. T-43. 29 pp.

Cihock, H. 2011. Oil leak shuts down Lake Bastrop. KXAN News, February 11, 2011. <http://www.kxan.com/dpp/news/local/oil-leak-shuts-down-lake-bastrop> Accessed July 12, 2011.

CH2M HILL. 2008. Climate change study report on evaluation methods and climate scenarios. Final draft report submitted to Lower Colorado River Authority and San Antonio Water System. Prepared by CH2M HILL, Austin, Texas. August 2008. 103 pp.

Cherry, D. S., J. L. Scheller, N. L. Cooper, and J. R. Bidwell. 2005. Potential effects of Asian clam (*Corbicula fluminea*) die-offs on native freshwater mussels (Unionidae) I: water-column ammonia levels and ammonia toxicity. *Journal of the North American Benthological Society* 24:369-380.

Christian, A.D., B.N. Smith, D.J. Berg, J.C. Smoot, and R.H. Findlay. 2004. Trophic position and potential food sources of 2 species of unionid bivalves (Mollusca: Unionidae) in 2 small Ohio streams. *Journal of the North American Benthological Society* 23:101-113.

City of San Antonio. 2010. Trends, challenges, and opportunities. Available at: [www.sanantonio.gov/planning/powerpoint/growth\\_trends\\_092506.pps](http://www.sanantonio.gov/planning/powerpoint/growth_trends_092506.pps) Accessed August 24, 2011.

Clary, K. H. 2010. Letter to PBS&J regarding LCRA Transmission Services Corporation (LCRA TSC) McCarney to Kendall to Gillespie transmission line facilities, competitive renewable energy zones (CREZ), Schleicher, Sutton, Menard, Kimble, Mason, Gillespie, Kerr, and Kendall Counties. April 1, 2010. 17 pp.

Clean Water Action. 2011. Conserving water in central Texas. <http://www.cleanwater.org/feature/conserving-water-central-texas> Accessed June 22, 2011.

Collier, M., R. Webb, and J. Schmidt. 1996. Dams and rivers: primer on the downstream effects of dams. U.S. Geological Survey Circular 1126. 94 pp.

Cooper, N. L., J. R. Bidwell, and D. S. Cherry. 2005. Potential effects of Asian clam (*Corbicula fluminea*) die-offs on native freshwater mussels (Unionidae) II: porewater ammonia. *Journal of the*

North American Benthological Society 24:381-394.

Couch, C. and P. Hamilton. 2002. Effects of urbanization on stream ecosystems. Fact Sheet FS-042-02. 2 pp.

Dall, W. H. 1882. American work on recent Molluca in 1881. The American Naturalist 16:953-968.

Delp, A. M. 2002. Flatworm predation on juvenile freshwater mussels. M. S. thesis, Southwest Missouri State University, Springfield, Missouri. 37 pp.

Dennis, S.D. 1984. Distributional analysis of the freshwater mussel fauna of the Tennessee River system, with special reference to possible limiting effects of siltation. PhD dissertation, Virginia Polytechnic Institute and State University, Blacksburg, VA. 247 pp.

Edwards, R. J. 1978. The effect of hypolimnion reservoir releases on fish distribution and species diversity. Transactions of the American Fisheries Society 107:71-77.

Ellis, M.M. 1936. Erosion silt as a factor in aquatic environments. Ecology 17:29 42.

Environmental Protection Agency (EPA). 2005. Aquatic life ambient water quality: diazinon. Final. EPA-822-R-05-006. Washington, DC. 85 pp.

EPA. 2007. National Water Quality Inventory: Report to Congress, 2002 Reporting Cycle. EPA 841-R-07-001. 39 pp.

Exelon. 2010. Victoria County Station early site permit application: Part 3 Environmental Report. Report to the Nuclear Regulatory Commission, Washington, D.C. 213 pp.

Forsage, A. and N. E. Carter. 1973. Effects of gravel dredging on the Brazos River. Proceedings of the Annual Conference of the Southeastern Association of Fish and Game Commissioners 27:695-709.

Fraley, S.J., and S.A. Ahlstedt. 2000. The recent decline of the native mussels (Unionidae) of Copper Creek, Scott County, Virginia. Pp. 189-195 in: P.D. Johnson and R.S. Butler, eds. Freshwater Mollusk Symposium Proceedings Part II: Proceedings of the First Symposium of the Freshwater Mollusk Conservation Society, March 1999, Chattanooga, Tennessee. Ohio Biological Survey, Columbus, Ohio.

Fuller, S. L. H. 1974. Clams and mussels (Mollusca: Bivalvia). Pp. 215-273 in Pollution Ecology of Freshwater Invertebrates. C. W. Hart and S. L. H. Fuller, eds. Academic Press, New York. 389 pp.

Garner, J. T., T. M. Haggerty, and R. F. Modlin. 1999. Reproductive cycle of *Quadrula metanevra* (Bivalvia: Unionidae) in the Pickwick Dam tailwater of the Tennessee River. American Midland Naturalist 141:277-283.

Gentner, H. W. and S. H. Hopkins. 1966. Changes in the trematode fauna of clams in the Little

Brazos River, Texas. *Journal of Parasitology* 52:458-461.

Gillespie County Soil and Water Conservation District 2011. Brush Clearing Programs. <http://www.gillespiecountyswcd.org/BrushClearing.html> Accessed June 15, 2011.

Gilpin, M. E., and M. E. Soule. 1986. Minimum viable populations: The processes of species extinctions. Pp 13-34 in *Conservation Biology: The Science of Scarcity and Diversity*, M.E. Soule, (ed.). Sinauer Associates, Sunderland, Mass.

Golladay, S. W., P. Gagnon, M. Kearns, J. M. Battle, and D. W. Hicks. 2004. Response of freshwater mussel assemblages (Bivalvia: Unionidae) to a record drought in the Gulf Coastal Plain of southwestern Georgia. *Journal of the North American Benthological Society* 23:494-506.

Gordon, M.E., and J.B. Layzer. 1989. Mussels (Bivalvia: Unionoidea) of the Cumberland River: review of life histories and ecological relationships. U.S. Fish and Wildlife Service Biological Report 89(15). 99 pp.

Goudreau, S., R. J. Neves, and R. J. Sheehan. 1993. Effects of wastewater treatment plant effluents on freshwater mollusks in the upper Clinch River, Virginia, U.S.A. *Hydrobiologia* 252:211-230.

Graf, D. L. and K. S. Cummings. 2007. Review of the systematics and global diversity of freshwater mussel species (Bivalvia: Unionoida). *Journal of Molluscan Studies* 73:291-314.

Greer, C. H. 2005. Hydrologic impacts of mechanical shearing of ashe juniper in Coryell County, Texas. Master's Thesis, Texas A&M University, San Antonio, Texas. 147 pp.

Groce, J. 2011. Email regarding Texas fatmucket in Onion Creek. Texas A&M University, San Antonio, Texas. June 6, 2011.

Guadalupe-Blanco River Authority. 2011. Canyon Reservoir. Available at: <http://www.gbra.org/canyon/default.aspx> Accessed June 13, 2011.

Haag, W. R. and A. M. Commens-Carson. 2008. Testing the assumption of annual shell ring deposition in freshwater mussels. *Canadian Journal of Fisheries and Aquatic Sciences* 65:493-508.

Hanson, J. M., W. C. Mackay, and E. E. Prepas. 1988. The effects of water depth and density on the growth of a unionid clam. *Freshwater Biology* 19:345-355.

Harrel, R. C. 1985. Effects of a crude oil spill on water quality and macrobenthos of a southeast Texas stream. *Hydrobiologia* 124:223-228.

Hartfield, P. W. 1993. Headcuts and their effect on freshwater mussels. Pp. 131-141 in: K.S. Cummings, A.C. Buchanan, and L.M. Koch, eds. *Conservation and Management of Freshwater Mussels*. Proceedings of a UMRCC Symposium, October 1992, St. Louis, Missouri. Upper Mississippi River Conservation Committee, Rock Island, Illinois.

- Hartfield, P. W. and E. Hartfield. 1996. Observations on the conglomerates of *Ptychobranhus greeni* (Conrad 1834) (Mollusca: Bivalvia: Unionoidea). *American Midland Naturalist* 135:370-375.
- Hastie, L. C., P. J. Cosgrove, N. Ellis, and M. J. Gaywood. 2003. The threat of climate change to freshwater pearl mussel populations. *Ambio* 32:40-46.
- Hersh, E. S. 2007. An integrated stream classification system for the state of Texas. *Surface Water Hydrology*, University of Texas, Austin. 43 pp. Available at: [https://webspace.utexas.edu/eh2489/CE394K2\\_Hersh\\_final.pdf](https://webspace.utexas.edu/eh2489/CE394K2_Hersh_final.pdf)
- Hickey, C. W., and M. L. Martin. 1999. Chronic toxicity of ammonia to the freshwater bivalve *Sphaerium novaezelandiae*. *Archives of Environmental Contaminants and Toxicology* 36:38-46.
- Horne, F. R. and S. McIntosh. 1979. Factors influencing distribution of mussels in the Blanco River of central Texas. *The Nautilus* 94:119-133.
- Howells, R. G. 1994. Distributional surveys of freshwater bivalves in Texas: progress report for 1992. *Texas Parks and Wildlife Management Data Series* 105. Austin, Texas. 20 pp.
- Howells, R. G. 1995. Distributional surveys of freshwater bivalves in Texas: progress report for 1993. *Texas Parks and Wildlife Management Data Series* 119. Austin, Texas. 50 pp.
- Howells, R. G. 1996. Distributional surveys of freshwater bivalves in Texas: progress report for 1994. *Texas Parks and Wildlife Management Data Series* 125. Austin, Texas. 45 pp.
- Howells, R. G. 1997a. Distributional surveys of freshwater bivalves in Texas: progress report for 1996. *Texas Parks and Wildlife Management Data Series* 144. Austin, Texas. 58 pp.
- Howells, R. G. 1997b. New fish hosts for nine freshwater mussels (Bivalvia:Unionidae) in Texas. *Texas Journal of Science* 49:255-258.
- Howells, R. G. 1998. Distributional surveys of freshwater bivalves in Texas: progress report for 1997. *Texas Parks and Wildlife Management Data Series* 147. Austin, Texas. 30 pp.
- Howells, R. G. 1999. Distributional surveys of freshwater bivalves in Texas: progress report for 1998. *Texas Parks and Wildlife Management Data Series* 161. Austin, Texas. 34 pp.
- Howells, R. G. 2000a. Distributional surveys of freshwater bivalves in Texas: progress report for 1999. *Texas Parks and Wildlife Management Data Series* 170. Austin, Texas. 56 pp.
- Howells, R. G. 2000b. Reproductive seasonality of freshwater mussels (Unionidae) in Texas. Pp. 35-48 in *Proceedings of the Conservation, Captive Care, and Propagation of Freshwater Mussels Symposium 1998*. Columbus, Ohio.
- Howells, R. G. 2001. Distributional surveys of freshwater bivalves in Texas: progress report for 2000. *Texas Parks and Wildlife Management Data Series* 187. Austin, Texas. 50 pp.

- Howells, R. G. 2002a. Distributional surveys of freshwater bivalves in Texas: progress report for 2001. Texas Parks and Wildlife Management Data Series 200. Austin, Texas. 28 pp.
- Howells, R. G. 2002b. Freshwater mussels (Unionidae) of the pimpleback complex (*Quadrula* spp.) in Texas. Texas Parks and Wildlife Management Data Series 197. Austin, Texas. 36 pp.
- Howells, R. G. 2003. Distributional surveys of freshwater bivalves in Texas: progress report for 2002. Texas Parks and Wildlife Management Data Series 214. Austin, Texas. 42 pp.
- Howells, R. G. 2004. Distributional surveys of freshwater bivalves in Texas: progress report for 2003. Texas Parks and Wildlife Management Data Series 222. Austin, Texas. 48 pp.
- Howells, R. G. 2005. Distributional surveys of freshwater bivalves in Texas: progress report for 2003. Texas Parks and Wildlife Management Data Series 233. Austin, Texas. 23 pp.
- Howells, R. G. 2006. Final report: statewide freshwater mussel survey. Federal Aid Grant number T-15-P. 106 pp.
- Howells, R. G. 2009. Biological opinion: conservation status of selected freshwater mussels in Texas. BioStudies, Kerrville, Texas. 25 pp.
- Howells, R. G. 2010a. Golden orb (*Quadrula aurea*): summary of selected biological and ecological data for Texas. BioStudies, Kerrville, Texas. 18 pp.
- Howells, R. G. 2010b. Smooth pimpleback (*Quadrula houstonensis*): summary of selected biological and ecological data for Texas. BioStudies, Kerrville, Texas. 18 pp.
- Howells, R. G. 2010c. Texas fatmucket *Lampsilis bracteata* (Gould 1855): summary of selected biological and ecological data for Texas. BioStudies, Kerrville, Texas. 20 pp.
- Howells, R. G. 2010d. Texas fawnsfoot (*Truncilla macrodon*): summary of selected biological and ecological data for Texas. BioStudies, Kerrville, Texas. 16 pp.
- Howells, R. G. 2010e. Texas pimpleback (*Quadrula petrina*): summary of selected biological and ecological data for Texas. BioStudies, Kerrville, Texas. 17 pp.
- Howells, R. G. 2010f. Freshwater mussels of Live Oak Creek, Gillespie County, Texas. Report for Settlers Ridge Homeowners and Kemp Smith, LLP. 18 pp.
- Howells, R.G. 2010g. Database of rare mussel collections in Texas, 1994 – 2004. Texas Parks and Wildlife Department Report generated May 2010 in excel spread sheet.
- Howells, R.G. 2012. Phone conversation regarding mussels within Guadalupe River. BioStudies. 8 August 2012.

- Howells, R. G., J. L. Dobie, W. L. Lindemann, and J. A. Crone. 2003. Discovery of a new population of endemic *Lampsilis bracteata* in central Texas, with comments on species status. *Ellipsaria* 5(2):5-6.
- Howells, R. G., R. W. Neck, and H. D. Murray. 1996. Freshwater mussels of Texas. Texas Parks and Wildlife Press, Austin, Texas. 218 pp.
- Hubbs, C., R. J. Edwards, and G. P. Garrett. 2008. An annotated checklist of the freshwater fishes of Texas, with keys to identification of species. Texas Academy of Science. 44 pp. Available at: <http://www.texasacademyofscience.org/>
- International Panel on Climate Change (IPCC). 2007. Summary for Policymakers. In: Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor and H. L. Miller (eds.). Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Jacobson, P. J., R. J. Neves, D. S. Cherry, and J. L. Farris. 1997. Sensitivity of glochidial stages of freshwater mussels (*Bivalvia*: Unionidae) to copper. *Environmental Toxicology and Chemistry* 16:2384-2392.
- Johnson, N.A. 2009. Database of freshwater mussels collected in Texas. University of Florida, Gainesville.
- Johnson, N.A., J. M. Pfeiffer III, P. D. Echo-Hawk, J. B. Moring, C. L. Stevens, and C. R. Randklev. 2014. Identification of Freshwater Mussels and Their Hosts in Texas Using DNA Barcodes. Power Point Presentation at Texas Freshwater Mussel Society Annual Workshop and Symposium 2014. Kerrville, Texas.
- Johnson, M. 2011. Email regarding Blanco River mussel surveys. Texas A&M University, San Antonio, Texas. July 9, 2011.
- Johnson, M.J. and J. Groce. 2011. Freshwater Mussel Surveys for Austin District of the Texas Department of Transportation (TxDOT). 10 pp.
- Joiner, A. 2010. Oil spill cleanup on Brazos River is continuing. Reporter-News, July 13, 2010. <http://www.reporternews.com/news/2010/jul/13/oil-spill-cleanup-on-brazos-river-is-continuing/?partne> Accessed July 12, 2011.
- Kanehl, P., and J. Lyons. 1992. Impacts of in-stream sand and gravel mining on stream habitat and fish communities, including a survey on the Big Rib River, Marathon County, Wisconsin. Wisconsin Department of Natural Resources Research Report 155. 32 pp.
- Karatayev, A. Y. and L. E. Burlakova. 2008. Final report: distributional survey and habitat utilization of freshwater mussels. Interagency final report to the Texas Water Development Board. Buffalo State College, Buffalo, New York. 47 pp.



- Karl, T.R., J.M. Melillo, and T.C. Peterson. 2009. Global climate change impacts in the United States. Cambridge University Press. 188 pp.
- Kat, P. W. and G. M. Davis. 1984. Molecular genetics of peripheral populations of Nova Scotioan Unionidae (Mollusca: Bivalvia). *Biological Journal of the Linnean Society* 22:157-185.
- Keen-Zebert, A. and J. C. Curran. 2009. Regional and local controls on the spatial distribution of bedrock reaches in the upper Guadalupe River, Texas. *Geomorphology* 112:295-305.
- Keller, A. E. and S. G. Zam. 1991. The acute toxicity of selected metals to the freshwater mussel *Anodonta imbecilis*. *Environmental Toxicology and Chemistry* 10:539-546.
- Kennon, F. W., J. T. Smith, and C. T. Welborn. 1967. Hydrologic studies of small watersheds: Escondido Creek, San Antonio River basin, Texas, 1955-63. Prepared for the Texas Water Development Board. 130 pp.
- Kolpin, D. W., E. T. Furlong, M. T. Meyer, E. M. Thurman, S. D. Zaugg, L. B. Barber, and H. T. Buxton. 2002. Pharmaceuticals, hormones, and other organic wastewater contaminants in U. S. streams, 1999-2000: a national reconnaissance. *Environmental Science and Technology* 36:1202-1211.
- Larralde, L. 2011. Interagency Initial Report to the San Antonio River Authority, A Longitudinal Survey and Habitat Utilization of Freshwater Mussels in the Lower San Antonio River. pp. 18.
- Layzer, J.B., M.E. Gordon, and R.M. Anderson. 1993. Mussels: the forgotten fauna of regulated rivers. A case study of the Caney Fork River. *Regulated Rivers: Research and Management* 8:63-71.
- Lee, M. C. and T. W. Schultz. 1994. Contaminants investigation of the Guadalupe and San Antonio River of Texas: 1992. U. S. Fish and Wildlife Service, Corpus Christi, Texas. 18 pp.
- Lehman, J. 2010. LCRA shuts down sand dredging operation. *The Llano News*, May 5, 2010. <http://www.llanonews.com/news/article/29725> Accessed June 20, 2011.
- Lewis, R., Jr. and F. L. Oliveria. 1979. Live oak decline in Texas. *Journal of Arboriculture* 5:241-244.
- Lewis, S. 2010. Oil spill in Brazos County prompts state inquiry. *The Eagle*, June 10, 2010. <http://www.theeagle.com/police/Oil-spill-in-Brazos-County-prompts-state-inquiry> Accessed July 12, 2011.
- Lower Colorado River Authority (LCRA). 2011a. LCRA dams form the Highland Lakes. Available at: <http://www.lcra.org/water/dams/index.html> Accessed June 13, 2011.
- LCRA. 2011b. Water quality permit review program: basin-wide permit review program.

<http://www.lcra.org/water/quality/protectingwaterqualitypage.html> Accessed June 20, 2011.

LCRA. 2011c. Texas drought: drought shows no signs of breaking. Drought update.

<http://www.lcra.org/water/drought/index.html> Accessed July 8, 2011.

MacCormack, Z. 2011. Kerrville fish kill blamed on sewage spill. MySanAntonio.com, April 26, 2011.

<http://www.mysanantonio.com/default/article/Kerrville-fish-kill-blamed-on-sewage-spill-1351985.php> accessed July 12, 2011.

Mace, R.E. and S.C. Wade. 2008. In hot water? How climate change may (or may not) affect groundwater resources of Texas. Gulf Coast Association of Geological Societies Transaction 58:655-668.

Magana, H. A. 2002. Invasive species emerging issues: toxic golden algae. U. S. Forest Service, Boise, Idaho. 2 pp.

Magnelia, S. J. 2007. Survival of rainbow trout fingerlings stocked into the special regulation zone of the Canyon Reservoir tailrace. Texas Parks and Wildlife Management Data Series 247. Austin, Texas. 32 pp.

March, F. A., F. J. Dwyer, T. Augspurger, C. G. Ingersoll, N. Wang, and C. A. Mebane. 2007. An evaluation of freshwater mussel toxicity data in the derivation of water quality guidance and standards for copper. Environmental Toxicology and Chemistry 26:2066-2074.

Marking, L. L. and T. D. Bills. 1979. Acute effects of silt and sand sedimentation on freshwater mussels. pp. 204-211 in: J.R. Rasmussen, ed. Proceedings of the UMRCC symposium on Upper Mississippi River bivalve mollusks. Upper Mississippi River Conservation Committee, Rock Island, Illinois.

Mashhood, F. 2011. Drought could dry Llano River by week's end, officials say. The Statesman, June 15, 2011.

<http://www.statesman.com/news/local/drought-could-dry-llano-river-by-weeks-end-1542491.html> accessed July 8, 2011.

May, M. 2011. Phone conversation regarding mussels in the Medina River, Texas. Texas Parks and Wildlife Department. June 9 2011.

May, M. 2012. Phone conversation regarding mussels in Pedernales River, Texas. Texas parks and Wildlife Department. April 20, 2011.

McKinney, M. L. 1997. Extinction vulnerability and selectivity: Combining ecological and paleontological views. Annual Review of Ecological Systems 28:495-516.

Meador, M. R. and A. O. Layher. 1998. Instream sand and gravel mining: environmental issues and regulatory process in the United States. Fisheries 23(11):6-13.

Minckley, W. L. and P. J. Unmack. 2000. Western springs, their faunas, and threats to their existence. Pp. 52-53 In: R.A. Abell, D.M. Olson, E. Dinerstein, P.T. Hurley et al. (eds). Freshwater Ecoregions of North America. Island Press, Washington, DC.

Mississippi Interstate Cooperative Resource Association (MICRA). 2005. Black carp risk assessment published. River Crossings 14(4):1-2.

Naimo, T. J. 1995. A review of the effects of heavy metals on freshwater mussels. *Ecotoxicology* 4:341-362.

National Response Center. 2010. Hazardous substance release/oil discharge notification affecting Keechi Creek in Leon County, Texas. 5 pp.

Neck, R. W. 1982a. Preliminary analysis of the ecological zoogeography of the freshwater mussels of Texas. Pp. 33-42 in J. R. Davis, ed. Proceedings of the Symposium of Recent Benthological Investigations in Texas and Adjacent States. Texas Academy of Science.

Neck, R. W. 1982b. A review of interactions between humans and freshwater mussels in Texas. Pp. 169 – 177 in J. R. Davis, ed. Proceedings of the Symposium of Recent Benthological Investigations in Texas and Adjacent States. Texas Academy of Science.

Neck, R. W. 1989. Freshwater bivalves of Medina Lake, Texas: factors producing a low-diversity fauna. *Texas Journal of Science* 41:319-325.

Neck, R. W. and R. G. Howells. 1994. Status survey of Texas heelsplitter, *Potamilus amphichaenus* (Frierson, 1898). Texas Parks and Wildlife Department Special Report. Austin, Texas. 51 pp.

Neves, R. J. 1987. Proceedings of the workshop on die-offs of freshwater mussels in the United States. Upper Mississippi River Conservation Committee, June 23-25, 1986, Davenport, Iowa. 14 pp.

Neves, R. J. 1991. Mollusks. Pp. 251-319 in: K. Terwilliger, coordinator. Virginia's endangered species. Proceedings of a symposium, April 1989, Blacksburg, Virginia. McDonald & Woodward Publishing Co., Blacksburg.

Neves, R. J., A. E. Bogan, J. D. Williams, S. A. Ahlstedt, and P. W. Hartfield. 1997. Status of aquatic mollusks in the southeastern United States: a downward spiral of diversity. Pp. 43-85 in: G.W. Benz and D.E. Collins, eds. Aquatic fauna in peril: the southeastern perspective, March-April 1994, Chattanooga, Tennessee. Special Publication 1, Southeast Aquatic Research Institute, Chattanooga.

Newton, T. J. 2003. The effects of ammonia on freshwater unionid mussels. *Environmental Toxicology and Chemistry* 22:2543-2544

Nichols, S.J. and D. Garling. 2000. Food-web dynamics and trophic-level interactions in a

multi-species community of freshwater unionids. *Canadian Journal of Zoology* 78:871-882.

Nico, L. G., and J. D. Williams. 1996. Risk assessment on black carp (Pisces: Cyprinidae). Unpublished report, U.S. Geological Survey, Gainesville, Florida. 61 pp.

Nielsen-Gammon, J. and B. McRoberts. 2009. An assessment of the meteorological severity of the 2008-09 Texas drought through July 2009. Office of the State Climatologist, College Station, Texas. 24 pp.

Noss, R. F. and A. Y. Cooperrider. 1994. Saving nature's legacy: Protecting and restoring biodiversity. Island Press, Washington, D.C.

Nueces River Authority. 2010. 2010 basin highlights report: San Antonio-Nueces coastal basin, Nueces River basin, Nueces-Rio Grande coastal basin. 44 pp.

Ohio State University Museum (OSUM). 2011a. Texas fatmucket (*Lampsilis bracteata*) records. Bivalve Database, Division of Molluscs, Museum of Biological Diversity, Department of Evolution, Ecology, and Organismal Biology. The Ohio State University, Columbus. Available at: <http://www.biosci.ohio-state.edu/~molluscs/OSUM2/> Accessed June 3, 2011.

OSUM. 2011b. Golden orb (*Quadrula aurea*) records. Bivalve Database, Division of Molluscs, Museum of Biological Diversity, Department of Evolution, Ecology, and Organismal Biology. The Ohio State University, Columbus. Available at: <http://www.biosci.ohio-state.edu/~molluscs/OSUM2/> Accessed June 3, 2011.

OSUM. 2011c. Smooth pimpleback (*Quadrula houstonensis*) records. Bivalve Database, Division of Molluscs, Museum of Biological Diversity, Department of Evolution, Ecology, and Organismal Biology. The Ohio State University, Columbus. Available at: <http://www.biosci.ohio-state.edu/~molluscs/OSUM2/> Accessed June 3, 2011.

OSUM. 2011d. Texas pimpleback (*Quadrula petrina*) records. Bivalve Database, Division of Molluscs, Museum of Biological Diversity, Department of Evolution, Ecology, and Organismal Biology. The Ohio State University, Columbus. Available at: <http://www.biosci.ohio-state.edu/~molluscs/OSUM2/> Accessed June 8, 2011.

OSUM. 2011e. Texas fawnsfoot (*Truncilla macrodon*) records. Bivalve Database, Division of Molluscs, Museum of Biological Diversity, Department of Evolution, Ecology, and Organismal Biology. The Ohio State University, Columbus. Available at: <http://www.biosci.ohio-state.edu/~molluscs/OSUM2/> Accessed June 9, 2011.

OSUM. 2011f. Mussel-host database. Division of Molluscs, Museum of Biological Diversity, Department of Evolution, Ecology, and Organismal Biology. The Ohio State University, Columbus. Available at <http://128.146.250.235/MusselHost/FMPro> Accessed June 9, 2011

Pappas, E. A., D. R. Smith, C. Huang, W. D. Shuster, and J. V. Bonta. 2008. Impervious surface impacts to runoff and sediment discharge under laboratory rainfall simulation. *Catena* 72:146-152.

- Peterjohn, W. T. and D. L. Correll. 1984. Nutrient dynamics in an agricultural watershed: observations on the role of a riparian forest. *Ecology* 65:1466-1475.
- Pringle, C. M. 1997. Exploring how disturbance is transmitted upstream: going against the flow. *Journal of the North American Benthological Society* 16:425-438.
- Pringle, C. M., M. C. Freeman, and B. J. Freeman. 2000. Regional effects of hydrologic alterations on riverine macrobiota in the new world: tropical-temperate comparisons. *Bioscience* 50:807-823.
- Raikow, D. F. and S. K. Hamilton. 2001. Bivalve diets in a midwestern U.S. stream. *Limnology and Oceanography* 46:514-522.
- Randklev, C. H. 2011a. San Saba River mussel collections. University of North Texas, Denton.
- Randklev, C. H. 2011b. *Quadrula houstonensis* and *Truncilla macrodon* localities. University of North Texas, Denton.
- Randklev, C.H. 2012. Annual report - scientific permit activities for permit # SPR-0511-142, report date 31 May 2011 through 22 May 2012.
- Randklev, C.H. 2012. Phone conversation regarding unpublished findings in multiple river basins. Institute of Renewable Natural Resources (IRNR). 1 October 2012.
- Randklev, C. H. and B. Lundeen. 2010. Comments regarding Texas fawnsfoot (*Truncilla macrodon*), smooth pimpleback (*Quadrula houstonensis*), and Louisiana pigtoe (*Pleurobema riddellii*) listings. Comments to U.S. Fish and Wildlife Service, Clear Lake, Texas. 10 pp.
- Randklev, C. R., B. J. Lundeen, R. G. Howells, and J. H. Kennedy. 2010a. First account of a living population of Texas fawnsfoot, *Truncilla macrodon* (Bivalvia: Unionidae), in the Brazos River, Texas. *The Southwestern Naturalist* 55:297-298.
- Randklev, C. R., B. Lundeen, and J. H. Kennedy. 2009. Final report: distributional survey and habitat utilization of freshwater mussels (Family Unionidae) in the lower Brazos and Sabine River basins. Interagency final report to the Texas Water Development Board. University of North Texas, Denton. 57 pp.
- Randklev, C. R., B. Lundeen, and J. H. Kennedy. 2010b. Summary of unpublished records for candidate mussel species from four museums in north central Texas. University of North Texas, Denton. 32 pp.
- Randklev, C. R., B. Lundeen, and J. H. Kennedy. 2010c. Unpublished museum records of rare freshwater mussels in Texas. University of North Texas, Denton.
- Ricciardi, A., R. J. Neves, and J. R. Rasmussen. 1998. Impending extinctions of North American freshwater mussels (Unionoida) following the zebra mussel (*Dreissena polymorpha*) invasion. *Journal of Animal Ecology* 67:613-619.
- Richter, B. D., D. P. Braun, M. A. Mendelson, and L. L. Master. 1997. *Conservation Biology*

11:1081-1093.

Robertson, C. 2011, Phone conversation discussing survey finding for Golden orb in 2010 for instream flows study. TPWD. 2010.

Roell, M. J. 1999. Sand and gravel mining in Missouri stream systems: aquatic resource effects and management alternatives. Missouri Department of Conservation, Columbia, Missouri. 26 pp.

Rogers, S. O., B. T. Watson, and R. J. Neves. 2001. Life history and population biology of the tan riffleshell (*Epioblasma florentina walkeri*) (Bivalvia: Unionidae). *Journal of the North American Benthological Society* 20:582-594.

San Antonio Water System. 2010. Collapsed pipe leads to sewer spill near San Antonio River. [http://www.saws.org/latest\\_news/Newsdrill.cfm?news\\_id=710](http://www.saws.org/latest_news/Newsdrill.cfm?news_id=710) accessed July 12, 2011.

Schneider, D. W., C. D. Ellis, and K. S. Cummings. 1998. A transportation model assessment of the risk to native mussel communities from zebra mussel spread. *Conservation Biology* 12:788-800.

Schnoor, J. L. and E. G. Fruh. 1979. Dissolved oxygen model of a short detention time reservoir with anaerobic hypolimnion. *Water Resources Bulletin* 15:506-518.

Serna, S. 2011. Big rig crashes into San Antonio River. KSAT News, May 10, 2011. <http://www.ksat.com/news/27838416/detail.html> accessed July 12, 2011.

Shaffer, M. L. 1981. Minimum population sizes for species conservation. *BioScience* 31:131- 134.

Shaffer, M.L., and F.B. Samson. 1985. Population size and extinction: a note on determining critical population size. *American Naturalist* 125:144-152.

Sharpe, A. J. 2005. What factors influence freshwater molluscan survival in the Conasauga River? M.S. thesis, North Carolina State University, Raleigh. 125 pp.

Shepard, W. D. 1993. Desert springs—both rare and endangered. *Aquatic Conservation: Marine and Freshwater Ecosystems* 3:351-359.

Silverman, H., S.J. Nichols, J.S. Cherry, E. Achberger, J.W. Lynn, and T.H. Dietz. 1997. Clearance of laboratory-cultured bacteria by freshwater bivalves: difference between lentic and lotic unionids. *Canadian Journal of Zoology* 75:1857-1866.

Simmang, C. M. and J. C. Curran. 2006. Morphological changes associated with gravel mining along the Colorado River, Texas. Texas State University, San Marcos, Texas.

Simpson, C.T. 1900. Descriptive catalog of the naiades, or pearly fresh-water mussels. *Proceedings of the United States National Museum* 22:501-1044. Available at: <http://books.google.com/ebooks?id=bRsraAAAYAAJ>

- Smith, D. G. 1985. Recent range expansion of the freshwater mussel *Anodonta imbecilis* and its relationship to clupeid fish restoration in the Connecticut River system. *Freshwater Invertebrate Biology* 4:105-108.
- Smith, R. L. 1974. *Ecology and Field Biology*. Second Edition. Harper & Row, New York, N.Y. 850 pp.
- Soil Conservation Service. 1959. Inventory and use of sedimentation data in Texas. Texas Board of Water Engineers Bulletin 5912. 94 pp.
- Sparks, B. L. and D. L. Strayer. 1998. Effects of low dissolved oxygen on juvenile *Elliptio complanata* (Bivalvia: Unionidae). *Journal of the North American Benthological Society* 17:129-134.
- Stanley, E. H., R. A. Short, J. W. Harrison, R. Hall, and R. C. Wiedenfeld. 1990. Variation in nutrient limitation in lotic and lentic algal communities in a Texas (USA) river. *Hydrobiologia* 206:61-71.
- Strayer, D. L. 1999. Effects of alien species on freshwater mollusks in North America. *Journal of the North American Benthological Society* 18:74-98.
- Strayer, D. L. 2008. *Freshwater mussel ecology: a multifactor approach to distribution and abundance*. University of California Press. 204 pp.
- Strayer, D. L., J. A. Downing, W. R. Haag, T. L. King, J. B. Layzer, T. J. Newton, and S. J. Nichols. 2004. Changing perspectives on pearly mussels, North America's most imperiled animals. *Bioscience* 54:429-439.
- Strecker, J. K. 1931. The distribution of the naiades or pearly fresh-water mussels of Texas. Baylor University Museum Special Publication Number 2. Waco, Texas. 71 pp.
- Texas Clean Rivers Program. 2008. Nitrate levels in the Concho River watershed. Available at [http://www.lcra.org/water/quality/crp/crpconcho\\_study.html](http://www.lcra.org/water/quality/crp/crpconcho_study.html) accessed August 24, 2011.
- Texas Clean Rivers Program. 2010a. Colorado River basin highlights report: water quality in the Texas Colorado River. 12 pp.
- Texas Clean Rivers Program. 2010b. Guadalupe River and Lavaca-Guadalupe coastal basins: basin highlights report. 48 pp.
- Texas Commission on Environmental Quality (TCEQ). 2010a. Basin 18: Guadalupe River. Available at [www.tnrc.state.tx.us/admin/topdoc/index.html](http://www.tnrc.state.tx.us/admin/topdoc/index.html)
- TCEQ. 2010b. Study of the methods for disposing of unused pharmaceuticals. Water Supply Division Report SFR-098. 278 pp.
- TCEQ. 2010c. Draft 2010 Texas 303(d) list. February 5, 2010. 106 pp.

Texas Natural Resources Conservation Commission. 2001. Implementation plan for Lake Austin dissolved oxygen TMDL for segment 1403. Austin, Texas. 12 pp.

Texas Parks and Wildlife Department (TPWD). 2002. Toxic golden algae in Texas. 23 pp.

TPWD. 2004. Sand, shell, gravel, and marl permit no. 2004-002 for Vulcan Construction Materials. Issued July 14, 2008.

TPWD. 2007a. Sand, shell, gravel, and marl permit no. 2007-1 for Whitley Dozer. Issued September 20, 2007.

TPWD. 2007b. General permit no. 2007-G14 for Cameron Fredkin. Issued August 28, 2007.

TPWD. 2008a. General permit no. 2008-G11 for Charles W. Evans. Issued April 23, 2008.

TPWD. 2008b. Sand, shell, gravel, and marl permit no. 2008-02 for Richmond Material Co. Issued November 3, 2008.

TPWD. 2008c. Sand, shell, gravel, and marl permit no. 2008-03 for the City of Austin. Issued December 1, 2008.

TPWD. 2009a. Lone zebra mussel found in Lake Texoma. News release.  
<http://www.tpwd.state.tx.us/newsmedia/releases/?req=20090421a> accessed July 8, 2011.

TPWD. 2009b. Sand and gravel general permit no. 2009-G 004 for Alan R. Stahlman. Issued March 9, 2009.

TPWD. 2009c. Zebra mussels spreading in Texas. News release.  
<http://www.tpwd.state.tx.us/newsmedia/releases/?req=20090817a> accessed June 22, 2011.

TPWD. 2010a. Historical fish kill events involving the golden alga, *Prymnesium parvum*, in Texas  
<http://www.tpwd.state.tx.us/landwater/water/environconcerns/hab/ga/blooms.phtml> accessed June 22, 2010.

TPWD. 2010b. Sand, shell, gravel, and marl permit no. 94-005D for Sand Supply/A Division of Campbell Concrete. Issued May 12, 2010.

TPWD. 2013. Zebra Mussels found in Lake Belton and suspected in Lakes Worth and Joe Pool. News Release. <http://www.tpwd.state.tx.us/newsmedia/releases/print.phtml?req=20130926a>, accessed September 26, 2013.

TPWD. 2014. Zebra Mussels Spread to Lake Waco. News Release. <https://tpwd.texas.gov/newsmedia/releases/?req=20141001b>, accessed October 1, 2014.

Texas Register. 2010. Chapter 65: Wildlife. Subchapter G. Threatened and endangered nongame



species. 31 TAC 65.175.

Texas Water Development Board (TWDB). 2011. Region K water plan for the Lower Colorado Regional Water Planning Group (adopted 2011). 130 pp. Available at [http://www.twdb.state.tx.us/wrpi/rwp/3rdRound/2011\\_RWP/RegionK](http://www.twdb.state.tx.us/wrpi/rwp/3rdRound/2011_RWP/RegionK).

Thomas, C. D. 1994. Extinction, colonization, and metapopulations: environmental tracking by rare species. *Conservation Biology* 8:373-378.

Turgeon, D.D., J.F. Quinn, Jr., A.E. Bogan, E.V. Coan, F.G. Hochberg, W.G. Lyons, P.M. Mikkelsen, R.J. Neves, C.F.E. Roper, G. Rosenberg, B. Roth, A. Scheltema, F.G. Thompson, M. Vecchione, and J.D. Williams. 1998. Common and scientific names of aquatic invertebrates from the United States and Canada: mollusks, 2nd edition. American Fisheries Society Special Publication 26, Bethesda, Maryland. 277 pp.

U. S. Army Corps of Engineers (USACE). 2010. Permit for Chemical Lime Company, Number SWF-2009-00317. Issued September 1, 2010.

U. S. Fish and Wildlife Service (Service). 2006. 5-year review of Higgins eye (*Lampsilis higginsii*). Bloomington, Minnesota. 25 pp.

U.S. Fish and Wildlife Service (Service). 2012. Field notes for presence/absence surveys conducted in 2012.

U.S. Fish and Wildlife Service (Service). 2013. Field notes for presence/absence surveys conducted in 2013.

U.S. Fish and Wildlife Service (Service). 2014. Field notes for presence/absence surveys conducted in 2014.

U. S. Geological Survey (USGS). 2001. Indications and potential sources of change in sand transport in the Brazos River, Texas. Water-Resources Investigations Report 01-4057. 38 pp.

USGS. 2011a. USGS 08138000 Colorado River at Winchell, TX.  
[http://waterdata.usgs.gov/tx/nwis/dv?cb\\_00060=on&cb\\_00010=on&cb\\_00095=on&cb\\_00065=on&for](http://waterdata.usgs.gov/tx/nwis/dv?cb_00060=on&cb_00010=on&cb_00095=on&cb_00065=on&for)  
Accessed June 22, 2011.

USGS. 2011b. USGS 08166200 Guadalupe River at Kerrville, TX.  
[http://waterdata.usgs.gov/tx/nwis/dv?cb\\_00060=on&cb\\_00065=on&format=gif\\_default&begin\\_date=2](http://waterdata.usgs.gov/tx/nwis/dv?cb_00060=on&cb_00065=on&format=gif_default&begin_date=2)  
Accessed June 24, 2011.

USGS. 2011c. Brazos River basin historical streamflow and water quality information.  
<http://www.brazos.org/HistoricalStreamFlowData.htm> Accessed June 27, 2011.

Valenti, T. W., D. S. Cherry, R. J. Neves, and J. Schmerfeld. 2005. Acute and chronic toxicity of mercury to early life stages of the rainbow mussel, *Villosa iris* (Bivalvia: Unionidae). *Environmental*

Toxicology and Chemistry 24:1242-1246.

Vannote, R. L., and G. W. Minshall. 1982. Fluvial processes and local lithology controlling abundance, structure, and composition of mussel beds. *Proceedings of the National Academy of Sciences* 79:4103-4107.

Vaughan, P. W. 1997. Winged mapleleaf mussel (*Quadrula fragosa*) recovery plan. U. S. Fish and Wildlife Service, Fort Snelling, Minnesota.

Vaughn, C. C. and C. M. Taylor. 1999. Impoundments and the decline of freshwater mussels: a case study of an extinction gradient. *Conservation Biology* 13:912-920.

Waller, D. L., J. J. Rach, and W. G. Cope. 1995. Effects of handling and aerial exposure on the survival of unionid mussels. *Journal of Freshwater Ecology* 10:199-208.

Wang, N., T. Augspurger, M. C. Barnhart, J. R. Bidwell, W. G. Cope, F. J. Dwyer, S. Geis, I. E. Greer, C. G. Ingersoll, C. M. Kane, T. W. May, R. J. Neves, T. J. Newton, A. D. Roberts, and D. W. Whites. 2007a. Intra- and interlaboratory variability in acute toxicity tests with glochidia and juveniles of freshwater mussels (Unionidae). *Environmental Toxicology and Chemistry* 26:2029-2035.

Wang, N., C. G. Ingersoll, D. K. Hardesty, C. D. Ivey, J. L. Kunz, T. W. May, F. J. Dwyer, A. D. Roberts, T. Augspurger, C. M. Kane, R. J. Neves, and M. C. Barnhart. 2007b. Acute toxicity of copper, ammonia, and chlorine to glochidia and juveniles of freshwater mussels (Unionidae). *Environmental Toxicology and Chemistry* 26:2036-2037.

Waters, T. F. 1995. Sediment in streams: sources, biological effects, and control. *American Fisheries Society Monograph* 7. 251 pp.

Watters, G. T. 1996. Small dams as barriers to freshwater mussels (*Bivalvia: Unionoida*) and their hosts. *Biological Conservation* 75:79-85.

Watters, G. T. 2000. Freshwater mollusks and water quality: effects of hydrologic and instream habitat alterations. Pp. 261-274 in: P. D. Johnson and R. S. Butler, eds. *Freshwater Mollusk Symposium Proceedings Part II: Proceedings of the First Symposium of the Freshwater Mollusk Conservation Society, March 1999, Chattanooga, Tennessee*. Ohio Biological Survey, Columbus.

Watters, G. T. and H. L. Dunn. 1995. The Unionidae of the lower Muskingum River (RM 34.1-0), Ohio, USA. *Walkerana* 7:224-263.

Watters, G. T. and S. H. O'Dee. 1999. Glochidia of the freshwater mussel *Lampsilis* overwintering on fish hosts. *Journal of Molluscan Studies* 65:453-459.

Watters, G. T. and S. H. O'Dee. 2000. Glochidial release as a function of water temperature: beyond bradycty and tachycty. Pp. 135-140 in R. A. Tankersly, D. I. Warmolts, G. T. Watters, and B. J. Armitage, eds. *Proceedings of the Conservation, Captive Care, and Propagation of*

Freshwater Mussels Symposium. Ohio Biological Survey, Columbus, Ohio. 274 pp.

Wilkins, Neal, J. Groce, and N. Ford. 2010. Freshwater mussel surveys within Travis County properties. November 2010. 10 pp.

Williams, J. D., S. L. H. Fuller, and R. Grace. 1992. Effects of impoundments on freshwater mussels (Mollusca: Bivalvia: Unionidae) in the main channel of the Black Warrior and Tombigbee Rivers in western Alabama. Bulletin of the Alabama Museum of Natural History 13:1-10.

Winemiller, K., N. K. Lujan, R. N. Wilkins, R. T. Snelgrove, A. M. Dube, K. L. Skow, and A. G. Snelgrove. 2010. Status of freshwater mussels in Texas. Texas A&M University, San Antonio. 64 pp.

Woodhouse, C.A., Meko, D.M., MacDonald, G.M., Stahle, D.W. and Cook, E.R. 2010. A 1,200-year perspective of 21st century drought in southwestern North America. Proceedings of the National Academy of Sciences USA 107: 21,283-21,288.

Yeager, M.M., D.S. Cherry, and R.J. Neves. 1994. Feeding and burrowing behaviors of juvenile rainbow mussels, *Villosa iris* (Bivalvia: Unionidae). Journal of the North American Benthological Society 13:217-222.

Young, W. C., D. H. Kent, and B. G. Whiteside. 1976. The influence of a deep storage reservoir on the species diversity of benthic macroinvertebrate communities of the Guadalupe River, Texas. Texas Journal of Science 27:213-224.

Zimmerman, L. L., R. J. Neves, and D. G. Smith. 2003. Control of predacious flatworms *Macrostomum* sp. in culturing juvenile freshwater mussels. North American Journal of Aquaculture 65:28-32.

## **Approval/Concurrence:**

Lead Regions must obtain written concurrence from all other Regions within the range of the species before recommending changes, including elevations or removals from candidate status and listing priority changes; the Regional Director must approve all such recommendations. The Director must concur on all resubmitted 12-month petition findings, additions or removal of species from candidate status, and listing priority changes.

Approve:



06/12/2015

Date

Concur:



12/15/2015

Date

Did not concur:

\_\_\_\_\_

\_\_\_\_\_

Date

Director's Remarks: